



# The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



## Brief communication

# Successful use of gentamycin as an antibiotic prophylaxis regimen to reduce the rate of healthcare-associated infections after renal transplantation

Cely S. Abboud\*, Maria Daniela Bergamasco, Ercilia Evangelista de Sousa, Eliana de Cássia Zandonadi, Doralice Cortez

Instituto Dante Pazzanese de Cardiologia, São Paulo, Brazil

### ARTICLE INFO

#### Article history:

Received 15 August 2012

Accepted 7 October 2012

Available online 28 February 2013

#### Keywords:

Renal transplantation

Antibiotic prophylaxis

Gentamycin

Health-care associated infections

### ABSTRACT

At our institution, we observed an increase in the incidence of healthcare-associated infections (HAI) due to Gram-negative bacilli, including three cases of carbapenem-resistant *Klebsiella pneumoniae*, among patients who underwent renal transplantation. In addition to strengthening infection control measures, we chose to add gentamycin to the antibiotic prophylaxis regimen of patients undergoing renal transplantation. We assessed the number of HAI occurring within 30 days of renal transplantation during two time periods: (1) the pre-intervention period, between September 2009 and June 2010, and (2) the post-intervention period, between July 2010 and April 2011. The intervention consisted of the addition of gentamycin to the surgical antibiotic prophylaxis regimen. The percentage of patients with HAIs was 31% lower during the post-intervention period ( $p=0.03$ ), with the greatest reductions observed for urinary tract infections ( $p=0.024$ ). Carbapenem-resistant *K. pneumoniae* was not isolated during this period. The investigated patients did not exhibit worsening renal function. Further studies are needed to assess antibiotic prophylaxis in renal transplantation patients at institutions where there is a high prevalence of multidrug-resistant Gram-negative bacteria.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/3.0/)

The optimal perioperative antibiotic prophylaxis regimen for renal transplant patients remains to be determined, and prophylaxis should be administered based exclusively on clinical evidence.<sup>1,2</sup>

At our institution, the Infection Control team observed an increase in the incidence of healthcare-associated infections (HAI) due to Gram-negative bacilli, including three cases of carbapenem-resistant *Klebsiella pneumoniae*, among patients

who underwent renal transplantation,<sup>3,4</sup> and measures were proposed to reduce the incidence of HAI.

The Instituto Dante Pazzanese de Cardiologia, located in São Paulo, is a public university hospital with a capacity of 350 beds for cardiovascular surgeries. Approximately 40 renal transplantation surgeries are performed annually. The Institution has a hospital infection control and prevention program and performs active disease surveillance, following criteria

\* Corresponding author at: Instituto Dante Pazzanese de Cardiologia, Av. Dr. Dante Pazzanese, 500, São Paulo, SP 04012-909, Brazil.

E-mail address: [cely.saad@gmail.com](mailto:cely.saad@gmail.com) (C.S. Abboud).

1413-8670 © 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/3.0/)

<http://dx.doi.org/10.1016/j.bjid.2012.10.016>

**Table 1 – Clinical characteristics of the patients during the pre- and post intervention periods.**

Characteristics of patients	Pre-intervention period	Post-intervention period	<i>p</i> *
N renal TX	24	21	
Average age	46.4	47.5	NS
DM 2	16.7% (4/24)	23.8% (5/21)	NS
Deceased donor	87.5% (21/24)	90.5% (19/21)	NS
Median length of surgery	300 min (5 h)	270 min (4 h 30)	
Double J	70.8% (17/24)	71.4% (15/21)	NS
Median use of drain	13 days	10 days	
Median of use IC	10 days	8 days	
Death within 30 days	8.3% (2/24)	4.8% (1/21)	NS
HAI rates within 30 days	54.2% (13/24)	19% (4/12)	0.03
Surgical site infection (SSI)	16/7% (4/24)	9.5% (2/21)	0.66
Urinary tract infection (UTI)	33.3% (8/24)	4.85 (1/21)	0.024
Primary bloodstream infection	4.2% (1/24)	0	
Pneumonia	0	4.8% (1/21)	

\* *p* value-Fischer exact test.

from the Centers for Disease Control and Prevention (CDC).<sup>5</sup>

We assessed the number of HAI occurring within 30 days of renal transplantation during two time periods: (1) the pre-intervention period, between September 2009 and June 2010, and (2) the post-intervention period, between July 2010 and April 2011.

The intervention consisted of an addition of gentamycin to the surgical antibiotic prophylaxis regimen. This addition was based on the previously determined sensitivity profile of HAI-causing bacteria. Antibiotic prophylaxis was administered during the post-intervention period. This regimen included two doses of gentamycin 2 mg/kg, one at the start of anesthesia and one 24 h later; one dose of cefuroxime 1.5 g at the start of anesthesia; and 750 mg of cefuroxime every 12 h for the next 48 h. Cefuroxime had been previously used at our institution.

The characteristics of the patients during the pre- and post-intervention periods are presented in Table 1.

The microorganisms isolated during the pre-intervention period were, 4 *Enterobacter* spp., 4 *Escherichia coli*, 1 *Proteus mirabilis*, 1 multi-sensitive *K. pneumoniae* and 3 carbapenem-resistant *K. pneumoniae*. In the post-intervention period 1 *Enterobacter cloacae*, 1 *E. coli* and 1 multi-sensitive *K. pneumoniae* were isolated.

The percentage of patients with HAIs was 31% lower during the post-intervention period ( $p=0.03$ ), with the greatest reductions observed for urinary tract infections ( $p=0.024$ ). Carbapenem-resistant *K. pneumoniae* was not isolated during this period.

The investigated patients did not exhibit worsening renal function.

The optimal regimen for perioperative antibiotic prophylaxis after renal transplantation remains to be determined. At our institution, we observed an increase in HAI in renal transplantation patients and, in addition to strengthening infection control measures we were encouraged to add gentamycin to the antibiotic prophylaxis regimen of patients undergoing renal transplantation. In the post-intervention period a significant reduction in HAI was observed ( $p=0.03$ ), with the greatest reductions observed for urinary tract infections ( $p=0.024$ ).

Based on these findings, gentamycin was included in the institutional protocol for surgical antibiotic prophylaxis for renal transplantation.

A limitation of this study is that only one public hospital in São Paulo was evaluated, such that the sample may not represent the general Brazilian population.

Further studies are needed to assess antibiotic prophylaxis in renal transplantation patients. These studies should include a larger number of patients and should be conducted at institutions where there is a high prevalence of multidrug-resistant Gram-negative bacteria.

## Conflict of interest

All authors declare to have no conflict of interest.

## Acknowledgement

We acknowledge Celia F. Haroguchi' contribution in execution of several steps in this study.

## REFERENCES

- Orlando G, Di Cocco P, D'Angelo M, et al. Surgical antibiotic prophylaxis after renal transplantation: time to reconsider. *Transplant Proc.* 2010;42:1118–9.
- Laftavi MR, Rostami R, Patel S, et al. Universal perioperative antimicrobial prophylaxis is not necessary in kidney transplantation. *Clin Transplant.* 2012;26:437–42.
- Bergamasco MD, Barroso Barbosa M, de Oliveira Garcia D, et al. Infection with *Klebsiella pneumoniae* carbapenemase (KPC)-producing *K. pneumoniae* in solid organ transplantation. *Transpl Infect Dis.* 2012;14:198–205.
- Abboud CS, Bergamasco MD, Doi A, et al. First Report of investigation into a outbreak due to carbapenemase-producing *Klebsiella pneumoniae* in a tertiary Brazilian hospital, with extension to a patient in the community. *J Infect Prev.* 2011;12:4.
- Garner JS, Jarvis WR, Emori TG, Horan TC, Hughes JM. CDC definitions for nosocomial infections, 1988. *Am J Infect Control.* 1988;16:128–40.