



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Letter to the Editor

Adherence to antiretroviral prophylaxis during early infancy in Latin America

Dear Editor,

We assessed reported adherence to antiretroviral (ARV) prophylaxis by HIV-exposed infants followed in the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD) International Site Development Initiative (NISDI) Longitudinal Study in Latin American Countries (LILAC) protocol¹ through at least 6–12 weeks of life. LILAC is a prospective cohort study at sites in Argentina, Brazil, and Peru which enrolled in 2008 and 2009. Among infants enrolled at birth, the primary caregiver was interviewed at birth and at every infant visit regarding infant adherence to ARVs during the previous three days, timing of last dose missed of any prescribed ARV, and any problems or situations making it difficult for caregivers to give infants the prescribed ARV(s).

Of the 385 infants prescribed ARVs at enrollment, percent adherence could be calculated for 312 infants (i.e., those with at least two days of available adherence data). Of these 312 infants, 309 (99.0%) had perfect adherence; 99% received zidovudine. Reasons cited for not having perfect adherence were: the maternity nurse did not administer the ARV(s) to the infant ($n=1$), the caregiver forgot to give the ARV(s) ($n=1$), and nausea ($n=1$). At the 6–12 week visit, 115 (30.6%) infants had a current prescription for ARVs. Percent adherence could be calculated for 97 infants at this visit, of whom 92 (94.8%) had perfect adherence. Caregivers of nine infants (8.0%) reported the last dose missed of any prescribed ARV was within the previous two weeks, six (5.3%) within the previous month, and two (1.8%) over a month ago. The remaining 96 caregivers (85.0%) reported no missed doses. Seventeen caregivers indicated difficulties in giving ARVs to the infants. Primary reasons cited were: the caregiver forgot to give the ARV(s) ($n=5$), the infant was sleeping ($n=2$), the caregiver ran out of medication ($n=2$), and the caregiver misunderstood the prescription ($n=2$). Given the high proportion of subjects reporting perfect adherence, formal analyses of factors associated with lack of adherence were not pursued.

Our results differ substantially from those of other studies in which lower adherence rates were reported from Africa^{2,3} (62–73.1% adherence to a single dose of nevirapine at birth)

and the U.S. (71% adherence to zidovudine in the week prior to interview)⁴ using similar self-report methods. It is unclear why higher adherence was observed in our population, although a higher susceptibility to social desirability bias is a possibility. Previous studies of adherence have demonstrated that self-reported adherence is generally higher than adherence measured by pill count, electronic monitoring, or drug concentrations in the blood.⁵ Thus, adherence rates in this cohort likely represent an overestimate of the actual adherence.

In conclusion, high levels of adherence to infant ARV prophylaxis were observed in this study. Infant ARV prophylaxis adherence should be further evaluated in studies utilizing additional measures of adherence in order to more precisely estimate adherence and understand factors predicting non-adherence.

Financial support

NICHD Contract # N01-HD-3-3345 (2002–2007) and by NICHD Contract # HHSN267200800001C (NICHD Control #: N01-HD-8-0001) (2007–2012).

Conflict of interest

All authors declare to have no conflict of interest.

Appendix A. The NISDI LILAC Study Team

Principal investigators, co-principal investigators, study coordinators, coordinating center representatives, and NICHD staff include: **Argentina:** Buenos Aires: Marcelo H. Losso, Irene Foradori, Alejandro Hakim, Erica Stankievich, Silvina Ivalo (Hospital General de Agudos José María Ramos Mejía); **Brazil:** Belo Horizonte: Jorge A. Pinto, Victor H. Melo, Fabiana Kakehasi, Beatriz M. Andrade (Universidade Federal de Minas Gerais); Caxias do Sul: Rosa Dea Sperhake, Nicole Golin, Sílvia Mariani Costamilan (Universidade de Caxias do Sul/Serviço Municipal de Infectologia); Nova Iguacu: Jose Pilotto, Luis Eduardo Fernandes, Gisely Falco (Hospital Geral Nova de Iguacu – HIV

Family Care Clinic); **Porto Alegre:** Rosa Dea Sperhacker, Breno Riegel Santos, Rita de Cassia Alves Lira (Universidade de Caxias do Sul/Hospital Conceição); Rosa Dea Sperhacker, Mario Ferreira Peixoto, Elizabete Teles (Universidade de Caxias do Sul/Hospital Fêmeina); Regis Kreitchmann, Luis Carlos Ribeiro, Fabrizio Motta, Debora Fernandes Coelho (Irmandade da Santa Casa de Misericórdia de Porto Alegre); **Ribeirão Preto:** Marisa M. Mussi-Pinhata, Geraldo Duarte, Adriana A. Tiraboschi Bárbaro, Conrado Milani Coutinho, Fabiana Rezende Amaral, Anderson Sanches de Melo (Hospital das Clínicas da Faculdade de Medicina de Ribeirão Preto da Universidade de São Paulo); **Rio de Janeiro:** Ricardo Hugo S. Oliveira, Elizabeth S. Machado, Maria C. Chermont Sapia (Instituto de Puericultura e Pediatria Martagão Gesteira); Esau Custodio Joao, Leon Claude Sidi, Maria Leticia Santos Cruz, Maria Isabel Gouvêa, Mariza Curto Saavedra, Clarisse Bressan, Fernanda Cavalcanti A. Jundi (Hospital dos Servidores do Estado); **São Paulo:** Regina Celia de Menezes Succi, Prescilla Chow (Escola Paulista de Medicina-Universidade Federal de São Paulo); **Peru:** Lima: Jorge O. Alarcón Villaverde (Instituto de Medicina Tropical “Daniel Alcides Carrión” – Sección de Epidemiología, UNMSM), Carlos Velásquez Vásquez (Instituto Nacional Materno Perinatal), César Gutiérrez Villafuerte (Instituto de Medicina Tropical “Daniel Alcides Carrión” – Sección de Epidemiología, UNMSM); **Data Management and Statistical Center:** Yolanda Bertucci, Rachel Cohen, Laura Freimanis Hance, René Gonin, D. Robert Harris, Roslyn Hennessey, James Korelitz, Margot Krauss, Sue Li, Karen Megazzini, Orlando Ortega, Sharon Sothorn de Sanchez, Sonia K. Stoszek, Qilu Yu (Westat, Rockville, MD, USA); **NICHD:** George K. Siberry, Rohan Hazra, Lynne M. Mofenson, Jennifer S. Read, Heather Watts (Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, Maryland, USA). Supported by NICHD Contract # N01-HD-3-3345 (2002–2007) and by NICHD Contract # HHSN267200800001C (NICHD Control #: N01-HD-8-0001) (2007–2012).

REFERENCES

1. Read JS, Duarte G, Freimanis Hance L, et al., for the NISDI Perinatal Study Group. The NICHD International Site Development Initiative perinatal cohorts (2002–09). *Int J Epidemiol.* 2012;41:642.
2. Kuonza LR, Tshuma CD, Shambira GN, Tshimanga M. Non-adherence to single dose nevirapine regimen for the prevention of mother-to-child transmission of HIV in Bindura town, Zimbabwe: a cross-sectional analytic study. *BMC Public Health.* 2010;10:218.
3. Barigve H, Levin J, Maher D, et al. Operational evaluation of a service for prevention of mother-to-child transmission of HIV

in rural Uganda: barriers to uptake of single-dose nevirapine and the role of birth reporting. *Trop Med Int Health.* 2010;15:1163–71.

4. Demas PA, Webber MP, Schoenbaum EE, et al. Maternal adherence to the zidovudine regimen for HIV-exposed infants to prevent HIV infection: a preliminary study. *Pediatrics.* 2002;110:e35.
5. Müller AD, Jaspan HB, Myer L, et al. Standard measures are inadequate to monitor pediatric adherence in a resource limited setting. *AIDS Behav.* 2011;15:422–31.

Cristina B. Hofer^{a,*}, D. Robert Harris^b, Mariza C. Saavedra^c, Jessica E. Haberer^d, Juliana Romeiro^e, Marisa M. Mussi-Pinhata^f, Erica Stankievich^g, Ivete M. Gomes^h, Regis Kreitchmannⁱ, Jennifer S. Read^j, for the NISDI LILAC Study Team[◇]

^a Instituto de Puericultura e Pediatria Martagão Gesteira and Department of Preventive Medicine, Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ, Brazil

^b Westat, Rockville, MD, USA

^c Hospital Federal dos Servidores do Estado, Rio de Janeiro, RJ, Brazil

^d Massachusetts General Hospital, Boston, MA, USA

^e Grupo de Estudos em HIV/AIDS, Medical School, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^f Department of Pediatrics, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, SP, Brazil

^g Hospital General de Agudos “J.M. Ramos Mejia”, Servicio de Inmunocomprometidos, Buenos Aires, Argentina

^h Hospital Geral de Nova Iguaçu, Rio de Janeiro, RJ, Brazil

ⁱ Irmandade da Santa Casa de Misericórdia de Porto Alegre, Porto Alegre, RS, Brazil

^j Pediatric, Adolescent, and Maternal AIDS Branch, Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, MD, USA

* Corresponding author at: Rua Bruno Lobo, 50, Rio de Janeiro, RJ, Brazil.

E-mail address: cbhofer@hucff.ufrj.br (C.B. Hofer).

◇ Members of the NISDI LILAC Study Team are shown in Appendix A.

Received 18 October 2012

Accepted 12 November 2012

Available online 13 May 2013

1413-8670

© 2013 Elsevier Editora Ltda.

Este é um artigo Open Access sob a licença de CC BY-NC-ND

<http://dx.doi.org/10.1016/j.bjid.2012.11.006>