



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

***Chlamydia trachomatis* in human immunodeficiency virus-infected men treated at a referral hospital for sexually transmitted diseases in the Amazonas, Brazil**[☆]

Alex Panizza Jalkh^{a,*}, Angelica Espinosa Miranda^{a,b}, Jose Camilo Hurtado-Guerreiro^a, Lorena Angelica Castano Ramos^a, Guisepppe Figliuolo^a, Jussimara Maia^a, Cintia Mara Costa^a, Rajendranath Ramasawmy^{a,c}, Luiz Carlos de Lima Ferreira^a

^a Fundação Medicina Tropical Doutor Heitor Vieira Dourado, Manaus, AM, Brazil

^b Infectious Diseases Unit, Federal University of Espírito Santo, Vitória, ES, Brazil

^c Universidade Nilton Lins, Manaus, AM, Brazil

ARTICLE INFO

Article history:

Received 28 February 2013

Accepted 19 June 2013

Available online 9 November 2013

Keywords:

Chlamydia trachomatis

Real-time polymerase chain reaction (RT-PCR)

HIV-infected men

Amazonas

ABSTRACT

Objectives: The aim of the present study was to determine the *Chlamydia trachomatis* prevalence and to identify the demographic, behavioural and clinical factors associated with *C. trachomatis* in human immunodeficiency virus infected men.

Study: This was a cross-sectional study of *C. trachomatis* prevalence among human immunodeficiency virus-infected men enrolled at the Outpatient clinic of acquired immunodeficiency syndrome of the Fundação de Medicina Tropical Dr. Heitor Vieira Dourado in Manaus, Amazonas, Brazil. *C. trachomatis* deoxyribonucleic acid from urethral samples was purified and submitted to real time polymerase chain reaction to identify the presence of *C. trachomatis*.

Results: A total of 276 human immunodeficiency virus-infected men were included in the study. The prevalence of *C. trachomatis* infection was 12% (95% confidence interval 8.1%–15.7%). The mean age of the participants was 34.63 (standard deviation 10.80) years. Of the 276 human immunodeficiency virus-infected men, 93 (56.2%) had more than one sexual partner in the past year and 105 (38.0%) reported having their first sexual intercourse under the age of 15 years. Men having sex with men and bisexuals amounted to 61.2% of the studied population. A total of 71.7% had received human immunodeficiency virus diagnosis in the last three years and 55.1% were using antiretroviral therapy. Factors associated with *C. trachomatis* infection in the logistic model were being single ($p < 0.034$), men having sex with men ($p < 0.021$), and having previous sexually transmitted diseases ($p < 0.001$).

Conclusion: The high prevalence of *C. trachomatis* infection among human immunodeficiency virus-infected men highlights that screening human immunodeficiency virus-infected men for *C. trachomatis*, especially among men having sex with men, is paramount to control the spread of *C. trachomatis* infection.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

[☆] Sources of support: Fundação de Medicina Tropical Dr. Heitor Vieira Dourado.

* Corresponding author at: Alameda Alaska 1051, ap. 601, Ed. Porto Seguro, Ponta Negra, Manaus, AM, Brazil.

E-mail addresses: ajalkh@gmail.com, alpajal@hotmail.com (A.P. Jalkh).

1413-8670 © 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de [CC BY-NC-ND](http://creativecommons.org/licenses/by-nc-nd/4.0/)

<http://dx.doi.org/10.1016/j.bjid.2013.06.007>

Introduction

Genital *Chlamydia trachomatis* (CT) infection, the most prevalent sexually transmitted disease (STDs) in the world, is a major public health problem.^{1,2} Asymptomatic infections are common and are associated with high probability of acquiring human immunodeficiency virus (HIV) infection.^{3–5} The screening of HIV-infected individuals for CT is a key strategy for reducing the transmission of HIV.^{2,3} Few studies have been conducted on CT infection among men in Brazil.⁶ A study in HIV-negative men attending STDs clinics have estimated the incidence of urethral CT infection to be 13.1%.⁷ Studies conducted in HIV-infected men outside of Brazil have reported prevalence rates of 9%⁸ and 2–10% in HIV-infected men and in men who have sex with men (MSM),^{2,9} respectively.

There is no mandatory reporting system for CT infections in Brazil,⁸ and the lack of population-based studies in HIV-infected patients hinders efforts to document the problem, implement priority interventions and evaluate their efficacy. Screening for CT infection in HIV-infected individuals is important for monitoring STDs and for elaborating preventive measures as well as treatment assistance to this population.

The present study aimed to determine the prevalence of CT infection and associated risk factors in HIV-infected men followed in a referral hospital specializing in infectious and tropical diseases in Manaus, Amazonas, Brazil.

Methods

A population based cross-sectional study among HIV-infected men was conducted at the referral hospital of the Fundação de Medicina Tropical Dr. Heitor Vieira Dourado (FMT-HVD) in Manaus, capital city of the state of the Amazonas, Brazil from November 1, 2009 to November 1, 2010. HIV-infected men were invited to participate in the study by signing an informed consent form as approved by the ethical committee of the FMT-HVD. Each participant answered a questionnaire concerning socio-demographic factors, age of first sexual intercourse, number of sexual partners over the past 12 months, regular use of condoms, sexual behaviours, prior symptoms of STDs. Signs and symptoms of STD on physical examination were also recorded and adequate treatment was provided.

The inclusion criteria were the following: male individuals with positive serology for HIV 1–2, sexually active, over 18 years of age and agreed to a urethral sample collection. Vulnerable male patients (mental diseases and younger than 18 years old) were excluded from the study.

The sample size was calculated for estimating the prevalence of CT infection in Human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS) people with a 95% confidence interval (CI) and a bilateral size of 0.5%. The sample size was calculated based on an average rate of 6.0% with a variation of $\pm 3.0\%$ which generated a number of 270 patients. Assuming a loss of 10%, the final sample size was 297 patients. Of note, the FMT-HVD is the referral centre for AIDS and all the patients with HIV/AIDS from the state

of Amazonas are referred to that centre. An average of 310 HIV-infected men is treated yearly from 2001 to 2010 at the FMT-HVD and in the year 2010, a total of 530 HIV-infected men were enrolled.

Urethral samples were collected by inserting a small brush (Cavi-brush) in the distal urethra, between 1.5 and 2 cm, and rotating it five times clockwise. The samples were then placed in tubes containing 400 μL of buffer solution (10 mM Tris-HCL and 1 mM EDTA) and kept on ice until they were transported to the laboratory, where they were stored at -20°C until processing for deoxyribonucleic acid (DNA) purification. DNA purification was performed using the commercial DNA extraction kit Nucleo-Spin Tissue – Machery – Nagel™ following the company's instructions. For the detection of CT DNA by RT-PCR, the following pair of primers amplifying a fragment of 241 base pairs CT plasmid DNA was used: KL1 5'-TCCGGAGCGAGTTACGAAGA-3' and KL 2'-AATCAATGCCCGGGATTGGT-3'. Briefly, 5 μL of DNA sample and 3 μL of each primer (10 pmol/ μL) were added to a final volume of 25 μL containing 12.5 μL of Maxima® SYBR Green/ROX qPCR Master Mix (2 \times). For the initial confirmation of the amplicon size the PCR product was electrophoresed in 1.5% agarose gel and revealed with ethidium bromide under UV. Furthermore, sample was considered positive when RT-PCR analysis showed unique peak dissociation curve at 78°C .

The data were stored and analyzed using the Epi Info software, version 3.5.2, and the Statistical Package for Social Science (SPSS) version 16.0 for Windows. Chi-square test with Yates correction was used to assess for possible associations between clinical, demographic or risk behaviours variables and CT infection. Fisher exact test was applied when appropriate. The odds ratio along with the 95% confidence interval (95% CI) was calculated for variables to estimate the degree of association between CT infection and potential risk factors. The multivariate logistic regression analysis was applied to examine the independent effect of each demographic, clinical or risk behaviour variables on CT infection, controlling for all other variables simultaneously. All variables that were moderately associated with a significance of $p \leq 0.15$ in the univariate analysis were considered for inclusion in the multivariate model. In the final model analysis, only those variables that remained significant with $p < 0.05$ were considered.

Results

Of a total of 276 HIV-infected men included in the study, 33 were infected with CT infection showing a prevalence of 12% (95% CI 8.1%–15.7%). No urethral symptoms were observed in any of the HIV-infected men. Baseline characteristics of the participants are shown in Table 1. The mean age of participants was 34.6 years (SD = 10.8) and 36.6% were 29 years old or less. More than 80% of the 276 HIV-infected men had up to 11 years of schooling. A total of 63.8% reported alcohol use and 23.9% illicit drugs abuse.

Concerning risk behaviours, as shown in Table 2, 93 (56.2%) participants reported having more than one sexual partner in the past year, and 105 (38.0%) had their first sexual intercourse under the age of 15 years. Prior to the diagnosis of HIV infection, 230 (83.3%) patients admitted irregular use

Table 1 – Prevalence of CT infection (RT-PCR) and its association with socio-demographic factors among 276 HIV-infected men treated at the FMT-HVD, 2009–2010.

Demographic Indices	n (%)	CT positive	CT negative	p-Value
<i>Age</i>				
18–29 years	101 (36.6)	2	22	0.3785 ^a
30–49 years	151 (54.7)	17	134	
≥50 years	24 (8.7)	2	22	
<i>Race/colour</i>				
Mulatto	210 (76.1)	21	189	0.066 ^a
White	48 (17.4)	11	37	
Black	18 (6.5)	1	17	
<i>Marital status</i>				
Single	158 (57.2)	25	133	0.034 ^a
Married/Living together	109 (39.5)	8	101	
Divorced	7 (2.5)	0	7	
Widower	2 (0.8)	0	2	
<i>Education</i>				
Primary	97 (35.1)	11	86	0.111 ^a
Secondary	125 (45.3)	11	114	
College	52 (18.8)	11	41	
Illiterate	2 (0.8)	0	2	
<i>Alcohol consumption</i>				
Yes	176 (63.8)	24	152	0.335 ^b
No	100 (36.2)	9	91	
<i>Use of illicit drugs</i>				
Yes	66 (23.9)	12	54	0.0837 ^b
No	210 (76.1)	21	189	

^a Pearson's chi-squared test (considered significant when $p \leq 0.05$).

^b Extended Fisher's exact test (considered significant when $p \leq 0.05$).

of condoms. After the diagnosis, 206 (74.6%) reported always using condoms. However, 179 (64.9%) subjects reported having previous STDs. A history of STDs was strongly associated ($p < 0.001$) with the detection of CT. Regarding sexual orientation, homosexuals and bisexuals together made up to 61.2% of the studied population. Multivariate logistic regression showed that MSM [OR=2.51 (95% CI 1.12–6.64) and previous STD [OR=3.46 (95% CI 1.96–5.72) were independent risk factors for CT infection among HIV-infected men.

Table 3 shows variables related to HIV status. None of the variables studied showed any association with CT infection. A total of 71.7% had received the HIV diagnosis in the last three years and 55.1% were using antiretroviral therapy.

Discussion

Our study prospectively investigated CT infection among HIV-infected men in the Amazonas through the RT-PCR technology to provide a rapid diagnosis for improving management of patients so as to limit the spread of CT. In Brazil, limited data are available demonstrating the precise pattern of the epidemiological behaviour of CT infection in men,^{6,10} and the status of this infection within the population of HIV-infected men is unknown.

The prevalence of CT urethral infection in HIV-infected men observed in this study was 12%. The observed prevalence is similar to that observed in non-HIV-infected men receiving

care at six STD clinics in Brazil,⁶ and in HIV-infected men in United Kingdom and USA.^{8,11}

STDs are of primary importance because of the emergence of AIDS. These infections enhance the sexual transmission of HIV^{3,12,13} and are associated with earlier and more severe symptoms in HIV-infected individuals.^{14–16} Furthermore, they can cause serious complications, resulting in chronic diseases and even death. Infections in men often cause problems in the genitourinary tract that can result in male infertility. In women, the consequences can be extremely severe, including chronic pelvic pain, infertility, and cervical cancer.^{16,17}

None of the HIV-infected men with CT infection in this study showed any urethral symptoms, which is consistent with most previous studies. A high proportion of asymptomatic patients has been observed in CT infections¹⁸ indicating a need for screening in this population.^{11,19–21} The routine testing of patients remains a controversial issue, given that some studies show a higher prevalence of CT infection among individuals with urethral symptoms.^{8,19,22–24}

Several studies have reported higher prevalence of CT infection in young individuals (<25 years of age),²³ but it was not observed in this study. Similar to work previously published by Iwuji et al. in 2008 in the United Kingdom,⁸ other studies have reported common associations between CT infections and reports of multiple partners, irregular use of condoms and a history of STDs.^{25,26} Here, only a history of STDs was found to be associated with CT infection. Notably, Fioravante et al.²⁵ observed a prevalence of 5% of chlamydial infection among male asymptomatic conscripts. This may

Table 2 – Sexual behaviours of 276 patients with HIV+/AIDS sorted by CT urethral infection status as detected by PCR.

Variables	n (%)	CT positive	CT negative	p-Value
Onset of sexual activity				
≤15 years	105 (38.0)	7	98	0.078 ^a
15–19 years	151 (54.7)	22	129	
≥20 years	20 (7.3)	4	16	
No. of sexual partners in the past year				
None	24 (8.7)	3	21	0.068 ^a
1	97 (35.1)	5	92	
2–9	93 (33.7)	14	79	
≥10	62 (22.5)	11	51	
Use of condoms before HIV+ diagnosis				
Sometimes	230 (83.3)	32	198	0.091 ^a
Always	7 (2.5)	0	7	
Never	39 (14.2)	1	38	
Use of condoms after HIV+ diagnosis				
No sex	34 (12.4)	5	29	0.766 ^a
Sometimes	29 (10.5)	4	25	
Always	206 (74.6)	24	182	
Never	7 (2.5)	0	7	
Sexual orientation				
MSM ^b	101 (36.6)	14	87	0.021 ^a
Bisexual	68 (24.6)	13	55	
Heterosexual	107 (38.8)	6	101	
Previous STDs				
Yes	179 (64.9)	30	149	0.001 ^a
No	97 (35.1)	3	94	

^a Pearson's chi-squared test (considered significant when $p \leq 0.05$).

^b Men who have sex with men.

Table 3 – Variables associated with the presence of HIV among 276 patients sorted by CT urethral infection status as detected by PCR.

Variables	n (%)	CT positive	CT negative	p-Value
Time since HIV diagnosis^a				
≤3 years	198 (71.7)	23	179	0.330 ^c
4–6 years	38 (13.8)	6	28	
7–9 years	15 (5.4)	0	15	
≥10 years	25 (9.1)	4	21	
Use of ART^b				
Yes	152 (55.1)	25	127	0.5778 ^d
No	124 (44.9)	8	116	
Period receiving ART^b				
<1 year	79 (52.0)	25	54	0.509 ^c
1–2 years	16 (10.5)	0	16	
2–3 years	12 (7.9)	0	12	
>3 years	45 (29.6)	8	37	
TCD4 Lymphocyte count (cells/mm³)				
<200	103 (37.3)	13	90	0.310 ^d
200–349	68 (24.6)	10	58	
350–500	41 (14.9)	4	37	
>500	40 (14.5)	6	34	
No results available	24 (8.7)	0	24	

^a Related to all 276 participants in the study.

^b Related to 152 patients who received antiretroviral therapy (ART).

^c Pearson's chi-squared test (considered significant when $p \leq 0.05$).

^d Extended Fisher's exact test (considered significant when $p \leq 0.05$).

reflect the prevalence in the general population of male Brazilians, which is lower than the prevalence in HIV-infected men.

A significant association between CT urethral infection and sexual orientation (MSM) was noted in this study. Several studies have demonstrated that a high prevalence of STDs,^{9,27} bacterial STDs,¹⁴ anorectal CT,^{22-24,28} and CT urethral infection^{14,28} are directly related to the sexual behaviour of MSM.

The possibility of response bias may be a limitation in this study due to the tendency to provide socially acceptable answers. However, such biases would result in underestimation of risk attitudes and behaviours. Inaccuracies of recall of condom use, age of first intercourse, and number of sexual partners may also have occurred. Another limitation is probably few HIV-infected individuals coming from high socioeconomic classes who may seek private clinics for treatment and thus potentially introducing patient selection bias. However, as treatment delivered for free in public hospitals in Brazil compared to the expensive cost in private hospitals, selection bias may have been negligible.

The findings from this study have important implications for education and prevention efforts directed towards young HIV-infected men, mainly MSM. A comprehensive approach would promote sexual responsibility and, at the same time, improve young men's understanding of sexual health risks to curb the spread of CT infection.

The lack of clinical symptoms in CT infection impairs the identification of infected individuals. This study underscores the need for screening through laboratorial tests to identify these individuals for rapid treatment, to limit the spread and also to avoid chronic symptoms and severe consequences for the individual. This topic is of extreme importance for health policies, as it has a significant impact on family policy and has consequences for the sexual and reproductive health of both the population in general and, more specifically, the HIV-infected population.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We acknowledge the contributions of Dra. Helene Silva Lira, Dra. Carolina Marinho da Costa, Dra. Marly Marquez de Melo and Technician Elizabeth Monteiro for performing laboratory procedures.

REFERENCES

- Centers for Disease Control and Prevention. Sexually transmitted diseases treatment, guidelines. *MMWR Morb Mortal Wkly Rep.* 2010;59:44-8.
- Centers for Disease Control and Prevention. Sexually transmitted diseases in the United States, 2008. National surveillance data for Chlamydia, gonorrhoea, and syphilis; 2010 <http://www.cdc.gov/std/stats08/trends.htm> [accessed 18.01.10].
- Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Infect.* 1999;75:3-17.
- Center for Diseases Control and Prevention. HIV prevention through early detection and treatment of other sexually transmitted diseases. United States. Recommendations of the Advisory Committee for HIV and STD Prevention. *MMWR Recomm Rep.* 1998;47:1-24.
- Brunham RC, Kimani J, Bwayo J, et al. The epidemiology of *Chlamydia trachomatis* within a sexually transmitted disease core group. *J Infect Dis.* 1996;173:950-6.
- Barbosa MJ, Moherdau F, Pinto VM, et al. *Gonorrhoea* and *Chlamydia* in men attending STD clinics. *Rev Soc Bras Med Trop.* 2010;43:500-3.
- Ministry of Health, Health Surveillance Secretariat, National Program of STD and AIDS. Prevalence and relative frequencies of sexually transmitted diseases (STD) in selected populations of Six Brazilian capitals, 2005. Brasília, DF: National Program of STD and AIDS; 2008.
- Iwuji CC, Reeves I, Nambiar K, et al. Diagnostic utility of urethral smears in predicting urethral *chlamydia* in HIV-infected men. *Int J STD AIDS.* 2008;19:741-3.
- Jin F, Prestage GP, Mao L, et al. Incidence and risk factors for urethral and anal *gonorrhoea* and *chlamydia* in a cohort of HIV-negative homosexual men: the Health in Men Study. *Sex Transm Infect.* 2007;83:113-9.
- Benzaken AS, Galban E, Moherdau F, et al. Prevalence of *Chlamydia trachomatis* infection and associated risk factors in different populations of both genders in Manaus city. *DST - J Bras Doencas Sex Transm.* 2008;20:18-23.
- Berry SA, Ghanem KG, Page KR, et al. Increased gonorrhoea and chlamydia testing did not increase case detection in an HIV clinical cohort 1999-2007. *Sex Transm Infect.* 2011;87(6):469-75.
- Joyee AC, Thyagarajan SP, Riddy EV, et al. Genital *Chlamydia* infection in patients: its relation to HIV infection. *Indian J Med Microbiol.* 2005;23:37-40.
- Lee HC, Ko NY, Chang CM, et al. Trends in sexually transmitted diseases and risk behaviors among HIV-infected patients at an outpatient clinic in southern Taiwan. *Sex Transm Dis.* 2010;37(2):86-93.
- Teague R, Mijch A, Fairley CA, et al. Testing rates for sexually transmitted infections among HIV-infected men who have sex with men attending two different HIV services. *Int J STD AIDS.* 2008;19:200-2.
- Wasserheit JN, Hillis SD. Screening for *Chlamydia* - a key to the prevention of pelvic inflammatory disease. *N Engl J Med.* 1996;334:1399-401.
- Brazil, Ministry of Health, Health Surveillance Secretariat, National Program of STD and AIDS. Sexually transmitted diseases control manual. Brasília: Ministry of Health; 2006.
- Stamm WE. *Chlamydia trachomatis* infections: progress and problems. *J Infect Dis.* 1999;179:S380-3.
- Paavonem J, Eggert-Kruse W. *Chlamydia trachomatis*: impact on human reproduction. *Hum Reprod Update.* 1999;5: 433-47.
- Johnson AS, Simms I, Sheringham J. The implementation of *chlamydia* screening: a cross-sectional study in the South East of England. *Sex Transm Infect.* 2010;86:217-21.
- Workowski KA, Berman SM, Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines. *MMWR Recomm Rep.* 2006;55:1-94.
- Schillinger JA, Dunne EF, Chapin JB, et al. Prevalence of *Chlamydia trachomatis* infection among men screened in 4 U.S. cities. *Sex Transm Dis.* 2005;32:74-7.
- Kent CK, Chaw JK, Wong W. Prevalence of rectal, urethral, and pharyngeal *chlamydia* and *gonorrhoea* detected in 2 clinical

- settings among men who have sex with men: San Francisco, California, 2003. *Clin Infect Dis.* 2005;41:67-74.
23. Annan NT, Sullivan AK, Nori A, et al. Rectal chlamydia – a reservoir of undiagnosed infection in men who have sex with men. *Sex Transm Infect.* 2009;85:176-9.
 24. Cook R, George St, Silvestre KA, et al. Prevalence of chlamydia and gonorrhoea among a population of men who have sex with men. *Sex Transm Infect.* 2002;78:190-3.
 25. Fioravante FC, Costa Alves MF, Guimarães EM, et al. Prevalence of *Chlamydia trachomatis* in asymptomatic Brazilian military conscripts. *Sex Transm Dis.* 2005;32:165-9.
 26. La Montagne DS, Fine DN, Marrazo JM. *Chlamydia trachomatis* infection in asymptomatic men. *Am J Prev Med.* 2003;24:36-42.
 27. Bachmann LH, Grimley DM, Waithaka Y, et al. Sexually transmitted disease/HIV transmission risk behaviours and sexually transmitted disease prevalence among HIV-positive men receiving continuing care. *Sex Transm Dis.* 2005;32:20-6.
 28. Ward H, Alexander S, Carder C, et al. The prevalence of lymphogranuloma venereum infection in men who have sex with men: results of a multicentre case finding study. *Sex Transm Infect.* 2009;85:173-5.