



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Arachidonic acid affects biofilm formation and PGE2 level in *Candida albicans* and non-albicans species in presence of subinhibitory concentration of fluconazole and terbinafine

Nripendra Nath Mishra ^{a,b}, Shakir Ali ^b, Praveen K. Shukla ^{a,*}

^a Medical Mycology Lab, Division of Fermentation Technology, CSIR-Central Drug Research Institute, B.S. 10/1, Sector 10, Jankipuram Extension, Sitapur Road, Lucknow 226021, India

^b Department of Biochemistry, Jamia Hamdard, New Delhi, India

ARTICLE INFO

Article history:

Received 26 May 2013

Accepted 13 September 2013

Available online 2 January 2014

Keywords:

Arachidonic acid

Candida albicans

AmBR

Candida tropicalis

Prostaglandin E2

FDA

ABSTRACT

Candida albicans utilizes arachidonic acid (AA) released during the course of infection (Candidiasis) from phospholipids of infected host cell membranes and synthesizes extracellular prostaglandin(s) which play an important role in hyphae formation and host cell damage. *C. albicans* biofilms secrete significantly more prostaglandin(s) and evidence suggests that *Candida* biofilms have dramatically reduced susceptibility to majority of antifungal drugs. AA influences the saturation level of lipids and fluidity of yeast cell membranes. Therefore the aim of this study was to evaluate the effect of AA alone or in combination with antifungal agents on biofilm formation and production of prostaglandin (PGE2) in *C. albicans*, *C. parapsilosis*, *C. glabrata*, *C. tropicalis*, and *C. albicans* amphotericin B resistant strain (AmBR). Maximum biofilm formation was found to be in the case of *C. albicans* compared to *C. non-albicans* species. However, among the non-albicans species *C. tropicalis* exhibited highest biofilm formation. Treatment with AA in combination with subinhibitory concentrations of fluconazole and terbinafine separately exhibited significant ($p < 0.05$) reduction in biofilm formation against *C. glabrata*, *C. parapsilosis*, *C. tropicalis* and AmBR as compared to their individual effect. Further, these two antifungal agents in combination with AA caused an increase in production of prostaglandin from fungal cell itself which was significant ($p < 0.05$) in case of all the strains tested.

© 2013 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND

Introduction

The pathogenic species of the genus *Candida* have gained recognition as nosocomial agents in recent years which may be attributed to specific risk factors associated with modern therapeutics like immunosuppressive agents, cytotoxic drugs,

and broad-spectrum antibiotics that suppress the normal bacterial microbiota. Likewise, use of various medical implants may result in *Candida* biofilm formation that often leads to aggravation of infection. Evidence suggests that *Candida* biofilms have dramatically reduced susceptibility to clinically used antifungal drugs like fluconazole and amphotericin B. Biofilm formation capacity of *C. albicans* and its role in

* Corresponding author.

E-mail address: pk_shukla@cdri.res.in (P.K. Shukla).

virulence and antifungal resistance has been studied in detail.¹ *Candida* non-albicans species like *C. parapsilosis* and *C. tropicalis* have also been found as biofilm forming pathogens in recent past.^{2,3}

Candida albicans has been reported to be responsible for the release of arachidonic acid (AA) from the host cells during infections^{4,5} which may modulate the cell growth, morphogenesis and invasiveness of causal agent by several modes. AA is a precursor for the production of eicosanoids which play an important role in morphogenesis and biofilm formation. Prostaglandin E₂ (PGE₂) is a primary product of AA metabolism in most of the eukaryotic cells that has also been reported in pathogenic fungi as well.^{6,7} Enhanced prostaglandin production during fungal infections could be one of the important factors in promoting colonization as well as chronic infections. The shift in host immune response toward increased colonization and chronic infections is due to PGE₂, which has ability to elicit both pro- and anti-inflammatory responses depending upon the host cells.^{8,9}

Exogenous AA has been reported to increase PGE₂ level significantly in *C. albicans* whereas the behavior of *Candida* non-albicans species and resistant strains in presence of AA is not much studied till now.⁸⁻¹⁰ The studies related to determining the level of PGE₂ in *C. albicans* and *C. non-albicans* species in presence of AA may help in understanding the biofilm forming capacity. The purpose of the present study was to compare the biofilm formation capacity and the level of prostaglandin secretion of the *C. albicans* and *Candida* non-albicans species in the presence of AA and known antifungal agents. The study also explores the pattern and amount of biofilm formation and the amount of prostaglandin secreted by a laboratory induced amphotericin B resistant *C. albicans* strain (AmBR).¹¹

Materials and methods

Candida strains used

Five species of *Candida* were used in this study. Out of these four species viz. *C. albicans* (ATCC-10231), *C. glabrata* (ATCC-MYA2950), *C. parapsilosis* (ATCC-22019), and *C. tropicalis* (ATCC-750) were purchased from ATCC (American type cell culture) while the fifth was an amphotericin B resistant strain of *C. albicans* (AmBR) developed earlier in our laboratory.¹¹

Media used and growth conditions

All the test strains maintained on Sabouraud dextrose agar were grown in yeast extract peptone and dextrose (YPD) medium (Difco BD, USA) for 24 h at 28 °C in an incubator shaker (180 rpm). The yeast cells were harvested, washed twice with 0.15 M phosphate-buffered saline (PBS, pH 7.4) and suspended in 10 ml of same buffer.

Antifungal susceptibility in vitro

The minimal inhibitory concentrations (MICs) of fluconazole (Sigma Aldrich, USA) and terbinafine (Biomole, USA) against all the five strains (as above) were determined by broth micro-dilution method as per guidelines of CLSI (formerly NCCLS)¹²

using RPMI 1640 (Sigma Aldrich, USA) buffered with MOPS (Sigma Chemical Co.) in 96-well tissue culture plates with positive and negative growth as well as drug controls. The test antifungals terbinafine and fluconazole were dissolved in DMSO (Merck, Germany) at a concentration of 2 mg/ml and 0.64 mg/ml respectively, diluted twofold and tested against 2×10^3 cells per ml of the test fungi. The micro-titer plates were incubated at 35 °C in a moist, dark chamber and MICs were recorded spectrophotometrically (Spectra Max, Molecular devices, USA) at 490 nm after 48 h incubation.

MICs of fluconazole and terbinafine were determined against biofilm formation by using the standard CLSI micro dilution protocol M27-A.¹²⁻¹⁴ Briefly, 1–5 $\times 10^3$ cells/ml were incubated in RPMI 1640 buffered with MOPS under shaking condition at 37 °C for 90 min and the plates were washed three times with PBS to remove non-adherent cells. The test antifungals fluconazole (32–0.25 µg/ml) and terbinafine (100–0.39 µg/ml) were diluted twofold in a replica plate and the whole contents transferred to the plates having the culture of biofilms and incubated for 48 h at 37 °C. Subsequently the plates containing *Candida* biofilms were washed thrice with PBS and 50 µl of XTT salt solution (1 mg/ml in PBS) and 4 µl of menadione solution (1 mM in acetone; Sigma) added to each well, incubated at 37 °C for 5 h, and XTT formazan measured in the supernatant medium at 490 nm by using a spectrophotometer (Molecular Devices, USA) where ≥80% decrease in formazan (OD at 490 nm) was taken as MIC. The experiment was done in duplicate and repeated three times on different occasions.

Biofilm formation

All the strains were subjected to biofilm formation under the following growth conditions separately: YNB with 50 mM glucose, YNB with 50 mM Glucose supplemented with 500 µM AA (Sigma Aldrich, USA), YNB with 50 mM glucose added subinhibitory concentration (1/2MIC) of fluconazole, YNB with 50 mM glucose added 1/2MIC terbinafine, YNB and 50 mM glucose supplemented with 1/2MIC fluconazole and AA and YNB supplemented with 50 mM glucose, 1/2MIC terbinafine and AA. Here it may be mentioned that subinhibitory concentration (1/2MIC) of both the drugs fluconazole and terbinafine against biofilm were used (Table 1). Biofilm formation was carried out in 96 well culture plates.¹⁵ Two-hundred µl fungal suspensions (1×10^6 cells) of all the test strains in the above medium were poured in 96 well tissue culture plates separately, kept under mild shaking condition for 90 min and subsequently the non-adherent cells were washed out with PBS. Then 200 µl media containing appropriate growth conditions as above was added in each well and incubated at 37 °C for 48 h.^{16,17}

Biofilm quantitation

Quantitation of *Candida* biofilms was performed by using 2,3-bis (2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino) carbonyl]-2H-tetrazolium hydroxide (XTT; Sigma Aldrich, USA) reduction assay where the water-soluble formazan product was measured at 490 nm.^{17,18} Following the biofilm formation phase, culture plates containing *Candida* biofilms

Table 1 – Minimal inhibitory concentration (MIC) of antifungal agents against different species and strains of *Candida*.

Strains	Terbinafine (MIC in µg/ml)				Fluconazole (MIC in µg/ml)			
	Planktonic		Biofilm		Planktonic		Biofilm	
	MIC	1/2MIC	MIC	1/2MIC	MIC	1/2MIC	MIC	1/2MIC
<i>C. albicans</i>	18.3	9.2	>100	100	0.36	0.18	4.0	2.0
AmBR	0.23	0.12	50.0	25.0	0.61	0.30	32.0	16.0
<i>C. glabrata</i>	72.0	36.0	>100	100	3.31	1.65	32.0	16.0
<i>C. parapsilosis</i>	0.29	0.14	50.0	25	0.74	0.37	2.0	1.0
<i>C. tropicalis</i>	68.0	34.0	>100	100	0.19	0.09	0.25	0.12

XTT reduction assay of biofilm.

were washed thrice with PBS and 50 µl of XTT salt solution (1 mg/ml in PBS) and 4 µl of menadione solution (1 mM in acetone; Sigma) added to each well. The plates were incubated at 37 °C for 5 h and XTT formazan measured in the supernatant medium at 490 nm by using a spectrophotometer (Molecular Devices, USA). The experiment was done in duplicate and repeated three times on different occasions.

Fluorescence microscopy

To study the biofilm formation fluorescence microscopy was performed in six well tissue culture plates and the images were obtained on Leica DM2500 Fluorescence microscope. Briefly, biofilms of above strains were grown in six well tissue culture plates in RPMI 1640 with L-glutamine and 0.165 M MOPS under different conditions of media as mentioned above. After incubation each well was washed with DPBS, stained with 100 mg/ml fluorescein diacetate (FDA; Sigma Aldrich, USA) and 50 mg/ml propidium iodide (PI; Sigma Aldrich, USA) for 3 h and viewed under microscope.¹⁹

Prostaglandin E₂ quantitation

Supernatant obtained (after 48 h) from the biofilm culture of all the strains under different growth conditions as mentioned above were subjected to PGE₂ estimation. Prostaglandin E₂ Express EIA kit (Cayman Chemicals) was used according to manufacturer's instructions. Briefly, cell culture supernatant was used directly for prostaglandin estimation and appropriate standards and controls were run simultaneously to obtain a standard curve. Each sample was assayed in triplicate and the experiments were repeated three times on different occasions.¹⁰ Sample concentrations were calculated on the basis of standard curve (plotted against standards versus PGE₂ concentration) and the data represented is the average of three experiments.

Statistical analysis

All experiments were performed in triplicate unless stated otherwise. The t-test was performed to determine the significance of the data sets.

Results and discussion

Candidiasis is one of the common fungal infections in humans and *C. albicans* being endogenous has been considered a

major causative agent but in recent past, but *Candida* non-albicans species have also marked their presence as frequent pathogens. The major clinical impact generated by *Candida* species is their ability to form biofilms which results in increased resistance to antifungal therapy and the ability of yeast cells within the biofilms to withstand host immune defences. Over and above it is interesting to note that there are species-specific variations in *Candida* biofilm morphology and formation capacity.^{1–3} Considering this, two antifungals, fluconazole and terbinafine were included in this study knowing the fact that the two are chemically different and yet block the synthesis of ergosterol.

Antifungal susceptibility against fluconazole and terbinafine

MICs were determined by CLSI method¹² for fluconazole and terbinafine against both planktonic and biofilm cells of *C. albicans*, *C. glabrata*, *C. parapsilosis*, *C. tropicalis* and AmBR. In the present study fluconazole (0.19–32 µg/ml) exhibited overall better spectrum of efficacy as compared to terbinafine (0.23–>100 µg/ml) against planktonic cells as well as biofilm cells (Table 1). However terbinafine exhibited an edge over fluconazole in case of AmBR and *C. parapsilosis* (Table 1).

Biofilm formation

In the present study, biofilm formation was studied in *C. albicans* and *Candida* non-albicans species (*C. glabrata*, *C. tropicalis*, and *C. parapsilosis*) and a laboratory developed amphotericin B resistant (AmBR) strain of *C. albicans*.¹¹ Polystyrene 96-well cell culture plates were used for biofilm formation in YNB media, which was measured by XTT reduction assay (Fig. 1). It is known that biofilms exhibit resistance against antifungals compared to planktonic cells; thus, the sub-inhibitory concentrations (1/2MIC, Table 1) of antifungal agents may have several important implications on planktonic cells destined to form biofilms.^{14,18,19} The influence of different treatments (growth conditions) was compared with the formation of biofilms in YNB alone which was considered as 100% for each of the five strains tested. In the present study YNB supplemented with AA (500 µM) did not affect the biofilm formation so the results indicate that 500 µM AA has no influence on the viability of the biofilms. The subinhibitory concentration of fluconazole caused reduced biofilm formation in case of *C. parapsilosis* (71.9% ± 2.9, *p* < 0.05) and the resistant strain AmBR

Table 2 – Biofilm formation (in %) under the influence of different media treatment.

Strains	% growth of biofilm under the influence of different media				
	YNB ^a +AA	YNB ^a +1/2FLZ	YNB ^a +1/2TER	YNB ^a +AA+1/2FLZ	YNB ^a +AA+1/2TER
<i>C. albicans</i>	95.9 ± 1.2	96.4 ± 1.02	93.6 ± 1.4	91.0 ± 2.4	86.1 ± 1.8
AmBR	94.1 ± 1.7	80.8 ± 3.2	47.2 ± 1.6	61.4 ± 1.9	26.5 ± 1.2
<i>C. glabrata</i>	93.6 ± 1.4	91.9 ± 2.9	62.3 ± 1.4	82.6 ± 2.3	31.6 ± 1.4
<i>C. parapsilosis</i>	90.3 ± 1.8	71.8 ± 2.9	42.7 ± 1.6	60.6 ± 2.2	23.0 ± 1.3
<i>C. tropicalis</i>	99.1 ± 1.1	96.6 ± 1.2	93.1 ± 1.4	88.6 ± 2.3	48.9 ± 2.4

^a Biofilm formation in YNB alone was considered as control (100%).

(81.9% ± 3.2, $p < 0.05$). The biofilms of *C. albicans* as well as *C. non-albicans* species are known to exert resistance against many antifungal agents including fluconazole,^{2,3} which has also been reported to reduce biofilm formation in clinically isolated fluconazole susceptible and resistant strains of *C. albicans*.¹⁷ However, in the present study a reduction in biofilm formation was evident in case of AmBR only (Fig. 1 and Table 2). The subinhibitory concentration of terbinafine caused significantly ($p < 0.0005$) reduced biofilm formation in case of *C. parapsilosis* (42.7% ± 1.6), AmBR (47.8% ± 1.6) and *C. glabrata* (61.2% ± 1.4) while in case of remaining two strains non-significant reduction was noticed ($p > 0.05$, Table 2).

It is encouraging to note that both the antifungals when tested in combination with AA resulted in further inhibition in biofilm formation as compared to either of the two agents (antifungal and AA) tested alone. Our results exhibit that fluconazole and terbinafine when tested in combination with AA caused least biofilm formation in case of *C. parapsilosis* followed by AmBR and *C. glabrata*. However, this effect was more pronounced ($p < 0.0005$) with terbinafine in *C. parapsilosis* (23% ± 1.3), AmBR (26.5% ± 1.2) and in case of *C. glabrata* it was


31.6% ± 1.4 (Fig. 1 and Table 2). In addition to this terbinafine with AA also reduced biofilm formation of *C. tropicalis* indicating that terbinafine with AA was the best combination in reducing biofilm formation against test strains as well as resistant strain AmBR (Table 2).

Fluorescence microscopy for biofilm evaluation

The biofilm cells of the test *Candida* strains were investigated by staining with FDA and PI and viewed under fluorescence microscope. The live and dead cells of biofilms were distinguished by green (FDA) and red (PI) fluorescence respectively indicating the presence of both viable and dead cells in varied amounts with regard to various treatments. The biofilm formation of all the test strains in YNB alone (Fig. 2A1-E1) were used as respective control where green fluorescent images were observed suggesting the viability of biofilm cells. Treatment of biofilms with various test agents like AA alone, 1/2MIC fluconazole, 1/2MIC terbinafine and 1/2MIC of antifungals combination with AA caused varied fluorescence (Fig. 2). The subinhibitory concentration (1/2MIC) of fluconazole and terbinafine also exhibited some changes (Fig. 2A3-E3 and A4-E4) in fluorescence as compared to their respective controls which is in agreement with the results of XTT assay (Fig. 1). As expected the subinhibitory concentration of fluconazole and terbinafine supplemented with AA resulted in significant suppression of biofilm formation (Fig. 2A5-E5 and A6-E6). Fluorescence microscopy of test strains with different treatments exhibited that 1/2MIC of antifungals in combination with AA affect the biofilm formation of all the test strains, which was greater in case of terbinafine as compared to fluconazole (Fig. 2).

PGE₂ production by biofilms of *C. albicans* and *C. non-albicans* strains

The planktonic and biofilm forms of *C. albicans* have been reported to produce eicosanoids particularly prostaglandin itself and/or in the presence of AA which is greater in biofilms.¹⁰ Prostaglandins have been known to be potential molecules to mediate cross talk between host and pathogen resulting in down-regulation of host immunological responses. Prostaglandins also play an important role in the persistence of fungal pathogen in immunocompetent hosts resulting in chronic infections. As stated above enhanced prostaglandin production has been observed in *C. albicans* biofilm¹⁰ whereas its production in biofilms produced by *Candida* non-albicans species and drug resistant strain has not

**Fig. 1 – XTT reduction assay of biofilm.**

Biofilm formation by *Candida albicans*, non-albicans, and AmBR under the influence of different media treatment was estimated by XTT assay. Biofilm formation in YNB alone was considered as control. Further 1/2MIC of fluconazole (FLZ) and terbinafine (TER) were supplemented in medium individually or in combination with arachidonic acid (AA). Results exhibit that FLZ and TER affects biofilm formation but it was more pronounced when tested in combination with AA in comparison to control ($p < 0.005$) and least biofilm formation was observed in case of *C. parapsilosis* followed by AmBR and *C. glabrata*. However, this effect was more pronounced with TER ($p < 0.005$) in comparison to FLZ.




Fig. 2 – Fluorescence microscopy of biofilm: fluorescence microscopy comparison of biofilms grown in presence and absence of arachidonic acid (AA) treated with subinhibitory concentration of fluconazole and terbinafine exhibited that least biofilm formation was observed in case of *C. glabrata* (Figs. B5 and B6), *C. parapsilosis* (Figs. C5 and C6) and AmBR (Figs. E5 and E6) compared to that in case of *C. albicans* (Figs. A5 and A6). The test strains with different treatments exhibited that 1/2MIC of antifungals with AA affect the biofilm formation of all test strains and among all treatments 1/2MIC of terbinafine with AA has greater effect than 1/2MIC fluconazole with AA. This result also suggests that terbinafine increases the susceptibility with AA as well as reduces the biofilm formation.

yet been studied. The amount of PGE₂ secreted by the test strains under control condition (YNB alone) for *C. albicans* (141 pg/ml, ± 6.2), *C. glabrata* (57.3 pg/ml, ± 1.3), *C. parapsilosis* (28.1 pg/ml, ± 1.1), *C. tropicalis* (29.6 pg/ml, ± 2.4) and AmBR (100.2 pg/ml, ± 2.1) was considered as basal concentration and the amount of PGE₂ was calculated after subtracting the basal concentration (Fig. 3). An increase in PGE₂ synthesis was noted for *C. albicans* (1387 ± 15.0 pg/ml), AmBR (1878 ± 11.0 pg/ml), *C. parapsilosis* (1189.3 ± 13.2 pg/ml), *C. glabrata* (1123.0 ± 15.5) and *C. tropicalis* (1846 ± 15.3 pg/ml) when grown in YNB

supplemented with AA ($p < 0.005$). These results suggest that AA could have been incorporated into the phospholipids of yeasts, influencing the saturation level and fluidity of yeast cell membranes. Influence of AA on cell membrane phospholipids unsaturation and ergosterol content has been reported recently.^{10,20} The reduced ergosterol content of amphotericin B resistant strain¹¹ could have been saturated with phospholipids incorporated by AA. Therefore increased amount of membrane phospholipids enables AmBR to produce more PGE₂ than its parent strain.

Table 3 – PGE₂ level (pg/ml) estimated under the influence of arachidonic acid (AA).

Strains	PGE ₂ level (pg/ml) in different conditions		
	YNB+AA ^a	YNB+AA+1/2MIC FLZ	YNB+AA+1/2MIC TER
<i>C. albicans</i>	1387 ± 15.0	2157.3 ± 26.1 ($p < 0.0001$)	1655.3 ± 14.1 ($p < 0.0001$)
AmBR	1878 ± 11.0	1935 ± 16.5 ($p < 0.005$)	2114 ± 12.2 ($p < 0.0001$)
<i>C. glabrata</i>	1123.0 ± 15.5	1174.3 ± 9.2 ($p < 0.05$)	1374.3 ± 16.5 ($p < 0.0001$)
<i>C. parapsilosis</i>	1189.3 ± 13.2	1944 ± 14.5 ($p < 0.0001$)	1942.3 ± 2.5 ($p < 0.0001$)
<i>C. tropicalis</i>	1846 ± 15.3	1966.6 ± 14.0 ($p < 0.005$)	1858 ± 9.1 ($p < 0.1$)

^a PGE₂ level in YNB+AA was used as basal concentration for comparison.

In this experiment the results show that prostaglandin production is enhanced when AA was supplemented with subinhibitory concentration of fluconazole, *C. albicans* exhibited 2157.3 ± 26.1 pg/ml PGE₂ followed by *C. tropicalis* (1966.6 ± 14.0 pg/ml), *C. parapsilosis* (1944 ± 14.5 pg/ml), AmBR (1935 ± 16.5 pg/ml) and *C. glabrata* (1174.3 ± 9.2 pg/ml) (Fig. 3 and Table 3). PGE₂ production in the presence of AA with subinhibitory concentration (1/2MIC) of terbinafine was more pronounced ($p < 0.0005$; Fig. 3 and Table 3) against AmBR (2114 ± 12.2 pg/ml) and *C. glabrata* (1374.3 ± 16.5 pg/ml). The present results open a new dimension in prostaglandin research as *Candida* non-albicans and *C. albicans* drug-resistant strain (AmBR) exhibited a notable difference in production of PGE₂ individually.

The variation in PGE₂ production in biofilms of the different test strains could be indicative of their persistence in host tissues. The level of PGE₂ in all test strains, including AmBR, was

found to be enhanced under medium of YNB with AA+1/2MIC fluconazole and with AA+1/2MIC terbinafine ($p < 0.0005$, Fig. 3). This can be explained by the fact that AA increases susceptibility of *C. albicans* as well as *Candida* non-albicans species against antifungal agents and may affect the prostaglandin production as well. Thus, it is evident from the results that the exposure to antifungal agents enhances the productions of PGE₂ to a greater extent in biofilms which may be responsible for development of resistance against antifungal agents.

Conclusion

The main general conclusion from this study is that the effect of subinhibitory concentrations of antifungal agents on biofilm formation is dependent on the type of antifungal agent and fungal strain used. Fluconazole and terbinafine when tested in combination with AA caused least biofilm formation in case of *C. parapsilosis* followed by AmBR and *C. glabrata*. However, this effect was more pronounced with terbinafine in terms of antifungal activity as well as reduced biofilm formation. Our results further suggest that AA could have been incorporated into the phospholipids of yeasts, influencing the saturation level and fluidity of yeast cell membranes. Further, the enhanced prostaglandin production in *Candida* non-albicans and drug resistant *C. albicans* as compared to *C. albicans* opens a new area of research to investigate the role of prostaglandin in antifungal resistance development and pathogenesis.




Fig. 3 – PGE₂ level estimation of *C. albicans* and non-albicans biofilms.

The level of PGE₂ produced by *Candida albicans* and non-albicans biofilm in different media treatment was estimated after 48 h incubation. Synthesis in YNB medium was used as control, further supplemented with arachidonic acid (AA), 1/2MIC of fluconazole (FLZ) and terbinafine (TER) individually. Result exhibited that PGE₂ formation was increased in the presence of 1/2MIC antifungals and it is more pronounced in *C. tropicalis* and AmBR against TER. The level of PGE₂ in all test strains including AmBR, was found to be enhanced under medium of YNB supplemented with AA, 1/2MIC FLZ ($p < 0.0005$) and AA, 1/2MIC TER ($p < 0.0005$). PGE₂ level was estimated by ELISA method (Cayman Kit). Result suggested that AA enhanced PGE₂ production in the presence of subinhibitory concentration of antifungals (1/2MIC).

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgement

We thank the Director of CDRI and head of the Fermentation Technology Division, CSIR-CDRI, Lucknow, India for providing facilities. We also thank ICMR, New Delhi for giving me fellowship during this work CSIR-CDRI Communication No; 8541.

REFERENCES

- Chandra J, Kuhn DM, Mukherjee PK, Hoyer LL, McCormick T, Mahmoud AG. Biofilm formation by the pathogen *C.*

- albicans*—development, architecture, and drug resistance. *J Bacteriol.* 2001;183:5385–94.
2. Sonia S, Mariana HA, Martins RO, et al. Biofilms of non-*Candida albicans* *Candida* species: quantification, structure and matrix composition. *Med Mycol.* 2009;47:681–9.
 3. Kuhn DM, Chandra J, Mukherjee PK, Ghannoum MA. Comparison of biofilms formed by *Candida albicans* and *Candida parapsilosis* on bioprosthetic surfaces. *Infect Immun.* 2002;70:878–88.
 4. Castro M, Ralston NV, Morgenthaler TI, Rohrbach MS, Limper AH. *Candida albicans* stimulates arachidonic acid liberation from alveolar macrophages through alpha-mannan and beta-glucan cell wall components. *Infect Immun.* 1994;62:3138–45.
 5. Deva R, Ciccoli R, Schewe T, Kock J, Nigam S. Arachidonic acid stimulates cell growth, and forms a novel oxygenated metabolite in *Candida albicans*. *Biochim Biophys Acta.* 2000;1486:299–311.
 6. Lamacka M, Sajbidor J. The occurrence of prostaglandins and related compounds in lower organisms. *Prostaglandins Leukot Essent Fatty Acids.* 1995;52:357–64.
 7. Lodewyk J, Kock J, Venter P, Botha A, et al. The production of biologically active eicosanoids by yeasts. *Adv Exp Med Biol.* 1997;433:217–9.
 8. Mairi C, Noverr Susan, Phare galen, Toews Michael, Coffey J, et al. Pathogenic yeasts *Cryptococcus neoformans* and *Candida albicans* produce immunomodulatory prostaglandins. *Infect Immun.* 2001;69:2957–63.
 9. Mairi X, Noverr C, Galen B, Toews, Gary B, Huffnagl. Production of prostaglandins and leukotrienes by pathogenic fungi. *Infect Immun.* 2002;70:400–2.
 10. Alem MAS, Duglas LJ. Prostaglandin production during growth of *Candida albicans* biofilms. *J Med Microbiol.* 2005;54:1001–5.
 11. Kumar R, Shukla PK. Amphotericin B resistance leads to enhanced proteinase and phospholipase activity and reduced germ tube formation in *Candida albicans*. *Fungal Biol.* 2010;114:189–97.
 12. National Committee for Clinical Laboratory and Standards. Reference method for broth dilution antifungal susceptibility testing of yeasts. Approved standard. NCCLS document M27-A. Wayne, PA: National Committee for Clinical Laboratory Standards; 1997.
 13. Ramage G, et al. Standardized method for in vitro antifungal susceptibility testing of *Candida albicans* biofilms. *Antimicrob Agents Chemother.* 2001;45:2475–9.
 14. Choi HW, Shin JH, Jung SI, Park KH, et al. Species-specific differences in the susceptibilities of biofilms formed by *Candida* bloodstream isolates to Echinocandin antifungals. *Antimicrob Agents Chemother.* 2007;51:1520–3.
 15. Shinde RB, Raut JS, Chauhan NM, Karuppayil SM. Chloroquine sensitizes biofilms of *Candida albicans* to antifungal azoles. *Braz J Infect Dis.* 2013;17:395–400.
 16. Bruzual I, Riggle P, Hadley S, Kumamoto CA. Biofilm formation by fluconazole-resistant *Candida albicans* strains is inhibited by fluconazole. *J Antimicrob Chemother.* 2007;59:441–50.
 17. Shin JH, et al. Biofilm production by isolates of *Candida* species recovered from nonneutropenic patients: comparison of bloodstream isolates with isolates from other sources. *J Clin Microbiol.* 2002;40:1244–8.
 18. Mohammed A, Al-Fattani, Douglas LJ. Biofilm matrix of *Candida albicans* and *Candida tropicalis*: chemical composition and role in drug resistance. *J Med Microbiol.* 2006;55:908–99.
 19. Nomanbhoy F, Steele C, Yano J, Yano J, Fidel Jr PL. Vaginal and oral epithelial cell anti-*Candida* activity. *Infect Immun.* 2002;70:7081–8.
 20. Ells R, Kock JL, Van Wyk PW, Botes PJ, Pohl CH. Arachidonic acid increases antifungal susceptibility of *Candida albicans* and *Candida dubliniensis*. *J Antimicrob Chemother.* 2009;63:124–8.