



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

The accessory gene regulator (agr) controls *Staphylococcus aureus* virulence in a murine intracranial abscesses model



Jian Gong^{a,d}, Dongzhi Li^{b,d}, Jun Yan^{c,d}, Yu Liu^c, Di Li^c, Jie Dong^c, Yaping Gao^c, Tao Sun^{b,*,**}, Guang Yang^{c,*,**}

^a Department of Neurosurgery, Beijing Tiantan Hospital, Capital Medical University, Beijing Neurosurgical Institute, Beijing, China

^b Ningxia Medical University, Incubation Base of National Key Laboratory for Cerebrocranial Diseases, Yinchuan, China

^c Beijing Institute of Basic Medical Sciences, Beijing, China

ARTICLE INFO

Article history:

Received 3 January 2014

Accepted 18 March 2014

Available online 13 May 2014

Keywords:

S. aureus

Agr

Intracranial abscesses model

ABSTRACT

Background: Intracranial abscesses are associated with high mortality. *Staphylococcus aureus* is one of the main pathogens that cause intracranial infection. Until now, there is no report to identify the key effectors of *S. aureus* during the intracranial infection.

Methods: The murine intracranial abscesses model induced by *S. aureus* was constructed. The vital sign and survival rate of mice were observed to evaluate the infection. Histological examination was used to diagnose the pathological alterations of mouse tissues. The sensitivity of *S. aureus* to whole blood was evaluated by whole-blood killing assay.

Results: In murine intracranial abscesses model, it was shown that the mortality caused by the accessory gene regulator (agr) locus deficient strain was significant decreased compared with its parent strain. Moreover, we found that RNAIII, the effector of agr system, was essential for the intracranial infection caused by *S. aureus*. In the further investigation, it was shown that restoration the expression of α -toxin in agr deficient strain could partially recover the mortality in the murine intracranial abscesses model.

Conclusion: Our data suggested that the agr system of *S. aureus* is an important virulence determinant in the induction and mortality of intracranial abscesses in mice.

© 2014 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND

Introduction

Intracranial abscesses frequently occur as a complication of sinusitis or neurosurgery, which are associated with high

mortality (19–43%).¹ Until now, the incidence, etiology, management and morbidity of intracranial abscesses are not well understood. *Staphylococcus aureus* is one of the main pathogens that cause intracranial abscesses.^{1–8}

* Corresponding author at: Institute of Basic Medical Sciences, 27 Taiping Road, Beijing 100850, China.

** Co-corresponding author.

E-mail addresses: suntao6699@163.com (T. Sun), yanggg@hotmail.com (G. Yang).

^d These authors contribute equally to this work.

<http://dx.doi.org/10.1016/j.bjid.2014.03.005>

1413-8670/© 2014 Elsevier Editora Ltda. Este é um artigo Open Access sob a licença de CC BY-NC-ND

Table 1 – Bacterial strains and plasmids.

Strain or plasmid	Comments	Source or reference
Strain		
<i>S. aureus</i>		
8325-4	Wild-type, rsbU ⁻	30
RN6390	Laboratory virulent strain derived from 8325	31
RN6911	RN6390 with an agr mutation, tet ^R	31
RN6911-hla	RN6911 restoring HLA activity, cmr ^R	This study
ΔRNAIII	Restriction-negative strain, 8325 derivative, kan ^R	32
ΔRNAIII-R	ΔRNAIII restoring RNAIII activity cmr ^R	32
<i>E. coli</i>		
DH5 α	A host strain for cloning	Transgene

The accessory gene regulator (agr) was identified as a quorum sensing system in *S. aureus*. RNAIII is the major effector of agr system. It acts as a small RNA regulating the expression of many virulence factors, including most of those encoding cell-wall-associated and extra-cellular proteins.^{9,10} Exotoxins expressed by *S. aureus* play important roles in the infection. α-Toxin is one of exotoxins, which is regulated by RNAIII. As a pore forming toxin, α-toxin can cause hemolysis and tissue damage. Recently, it has been found that α-toxin can also interact with its receptor ADAM10 to aggravate *S. aureus* infection.¹¹

Herein we describe an animal model of intracranial abscesses caused by *S. aureus* that we have constructed. The pathogenic role of agr system is investigated in this model.

Materials and methods

Mice

Adult female C57BL/6J mice (6–8 weeks old), obtained from Vital River Laboratories, were used for all experiments. Mice were housed in group cages, maintained on a 12:12 hour light-dark cycle, in a controlled environment (25 °C) and given unrestricted access to food and water. All animal experimental protocols of the study are in accordance with the national guidelines for the use of animals in scientific research “Regulations for the Administration of Affairs Concerning Experimental Animals”. It is also approved by Animal Care and Use Committee of Beijing Institute of Basic Medical Sciences with the approval number BMS-130112.

Staphylococcus aureus strains and culture conditions

S. aureus cells were grown in 5 mL of brain heart infusion (BHI), and *Escherichia coli* cells were grown in Luria-Bertani (LB) medium. Bacteria were cultured at 37 °C for 12 h with shaking at 200 rpm in a 25-mL test tube. Cells were transferred from 1 mL of pre-culture to 100 mL of BHI or LB medium in a 500-mL flask. *S. aureus* cells were routinely grown in BHI and *E. coli* cells were grown in LB medium either with no antibiotics, or with 40 µg/mL Chloramphenicol, 50 µg/mL tetracycline, 100 µg/mL ampicillin, and 50 µg/mL kanamycin.

The strains used in this study are listed in Table 1.

Mouse brain infection model

Mice were anesthetized with 1% pentobarbital sodium (3–5 mg/100 gbw i.p.). While under anesthesia, mice were prepared for injection by securing the head in a stereotactic frame and quickly locating appropriate coordinates in the frontal cortex region on the skull. The general location for injection is 1 mm anterior to bregma and 1 mm lateral to the sagittal suture on the right side.¹² Parting the fur with a needlepoint along both axes is appropriate for injection. To make an injection, orient the needle perpendicular to the skull and press through the calvarium to approximately 2 mm depth. Inject 5 µL bead suspension (10⁶ CFU) into the brain by pressing the syringe button down and waiting 2 s for the bacteria to enter the brain. Control mice received injections of sterile saline. Carefully remove the needle and place the mouse in a clean, warm cage. All mice survived the injection procedure. After recovery from anesthesia, mice were kept at room temperature (25 °C) and provided with standard chow and watered *ad libitum*. 48-hour mortality, average time of death and rectal temperature were monitored.

Histological examination of mouse tissues

Mice were killed 12 h after injection and brains were removed immediately. Mouse brain was fixed in 10% neutral-buffered formalin for 48 h. The fixed tissue was processed through graded series of ethanol, followed by xylene, and embedded in paraffin. Tissue blocks through the entire abscess were sectioned at 5 µm and slides stained with hematoxylin-eosin.

Whole-blood killing assays

Bacteria cultures were washed twice in PBS, diluted to an inoculum of 10⁵ CFU in 25 µL PBS, and mixed with 75 µL of freshly drawn mouse blood in heparinized tubes. The tubes were incubated at 37 °C for 4 h with agitation, at which time dilutions were plated on BHI agar for enumeration of surviving CFU.¹³




Fig. 1 – Construction of murine intracranial abscesses model caused by *S. aureus*. (A) Measurement of rectal temperature of mice at different time points after challenge with *S. aureus* 8325-4 (0 h, 5 h, 7.5 h, 10 h, 12.5 h, 15 h) ($n = 9$ in each group). (B) Measurement of survival curve of mice in the two groups ($n = 52$ mice in each group). Mice were challenged with *S. aureus* 8325-4 or saline. The number of dead mice was recorded. The survival rates of the two groups were calculated individually, $p < 0.0001$. (C) Histological examination of mouse brain tissue. Brain tissues from the two groups were collected 8 h post-challenge with *S. aureus* 8325-4 and fixed. The fixed tissues were stained by H&E. (D) White blood cell count of the two groups at different time points (0 h, 4 h and 8 h) post-challenge with *S. aureus* ($n = 8$ in each group). (E) Percentage of neutrophils in the two groups at different time points (0 h, 4 h and 8 h) post-challenge with *S. aureus* ($n = 8$ in each group).

Restoring HLA activity in RN6911

The whole HLA coding region was amplified by PCR and cloned into pOS1 vector. The resulting plasmid (pOS1-hla) was transformed into *E. coli* DH5 α . Positive clones were selected on LB plates containing 100 μ g/mL ampicillin. The recombinant plasmid was isolated from positive clones and transformed to *S. aureus* RN4220 cells. Transformants were selected on tryptic soy agar plates containing 40 μ g/mL chloramphenicol at 37 °C. The positive clone was selected by colony PCR. Then the plasmid was isolated and transformed to RN6911. The transformants were selected on tryptic soy agar plates containing 40 μ g/mL chloramphenicol at 37 °C for 20 h. Plasmids of positive restoring colonies were extracted and confirmed by PCR.

Statistical analysis

Data are expressed as mean \pm SEM. The significance of differences between experimental groups was determined by two-tailed t-test assuming a 95% confidence interval.

Results

Construction of murine intracranial abscesses model caused by *S. aureus*

Injection of *S. aureus* 8325-4 into murine frontal cortex could produce a focal brain abscess with a clinical course similar to that seen in clinical patients with intracranial infection. Within 48 h post bacterial challenge, mice demonstrated clinical signs of illness including hunched posture, ruffled fur, lethargy, spasm, and finally expired. In contrast, any overt changes in posture, grooming, or physical activity of mice injected with sterile saline were not observed. However, challenge with *S. aureus* could not induce significant changes of rectal temperature (Fig. 1A). More than 50% of mice were dead after 48 h (Fig. 1B). Brains were collected 8 h after injection of Bacteria. A well-circumscribed abscess containing neutrophil and macrophage infiltrate, with significant mass effect was observed in histological examination of brains injected with *S. aureus* (Fig. 1C). Meanwhile, we also evaluate the hemogram




Fig. 2 – The role of agr system in murine intracranial abscesses model. (A) Survive curve of mice challenged with different strains ($n = 16$ mice in each group). Mice were challenged with RN6390B, RN6911, and saline, respectively. The number of dead mice was recorded. The survival rate of the two groups was calculated individually. **(B)** Histological examination of mouse brain tissue. Brain tissues from RN6390B and RN6911 groups were collected at 12 h post-challenge with bacteria and fixed. The fixed tissues were stained by H&E. **(C)** Percentage of neutrophils in the three groups (RN6390B, RN6911 and saline) at different time points (0h, 4h and 8h) post-challenge with *S. aureus* ($n = 10$ in each group). **(D)** Survive curve of mice challenged with different strains ($n = 22$ mice in each group). Mice were challenged with *S. aureus* 8325-4, Δ RNAIII, Δ RNAIII-R, and saline, respectively. The number of dead mice was recorded. The survival rate of the two groups was calculated individually. **(E)** Survival percentage of the different strains (RN6390B and RN6911) in murine whole blood. Bacteria (1×10^5 CFU) were incubated with murine whole blood for 4 h. Survival rate of bacteria were determined at BHI plate.

of mice at 4 h and 8 h after infection. The number of white blood cells was not significantly altered (Fig. 1D), but the percentage of neutrophil was significantly increased at different time points in this model (Fig. 1E).

The role of agr system in murine intracranial abscesses model

Given that agr system can regulate most of exotoxins expressed by *S. aureus*, we tried to determine whether agr system was involved in intracranial infection. Mice were injected with *S. aureus* RN6390B or RN6911 (agr-null strain). Injection with RN6911 could not induce death of mice (Fig. 2A). However, histological examination showed that RN6911 induced similar histological alteration of brain compared to RN6390B (Fig. 2B). The percentage of neutrophil was significantly increased in RN6390B and RN6911 groups, compared to sterile saline (Fig. 2C). It is well known that RNAIII is the key

effector of agr system. Further investigating, we determined whether RNAIII deletion as well as agr-null strain was associated with decreased mortality of mice. The RNAIII deletion strains (Δ RNAIII) and the restored strain (Δ RNAIII-R) were previously constructed. Mice were challenged with wild-type *S. aureus* 8325-4, Δ RNAIII and Δ RNAIII-R, respectively. In line with expectation, Δ RNAIII could not induce the death of mice and challenge with the restoration RNAIII strain recovered virulence and the consequent mortality of mice (Fig. 2D). All these results suggested that agr system was essential for *S. aureus* induced mortality in the murine intracranial abscesses model.

As the alteration of hemogram induced by RN6911 was not different from that of RN6390B, we wondered if a deficient agr system could make the bacteria more sensitive to whole blood killing. RN6911 and RN6390B were incubated with murine whole blood respectively. The bacterial survival rate was assessed 4 h after incubation. The survival rate of RN6911




Fig. 3 – Determination of the role of α -toxin in murine intracranial abscesses model. Survive curve of mice challenged with different strains ($n = 12$ mice in each group). Mice were challenged with RN6390B, RN6911, RN6911-hla, and saline, respectively. The number of dead mice was recorded. The survival rate of the two groups was calculated individually.

bacteria was not significantly different when compared to RN6390B bacteria (Fig. 2E). This result suggests that the mortality alteration caused by agr null strain was not related with the sensitivity to whole blood killing.

α -Toxin was involved in the mortality caused by *S. aureus* in murine intracranial abscesses model

Exotoxins expressed by *S. aureus* are important for the infection.^{10,14,15} Some exotoxins are regulated by the agr system^{10,16} and α -toxin is one of the major exotoxins regulated by agr system. RN6911 do not express α -toxin. In the following study, we investigated whether α -toxin was involved in intracranial abscesses. The expression of α -toxin was recovered in RN6911 by the plasmid pOS1-HLA to generate strain (RN6911-HLA). Mice were challenged with three different strains (RN6390B, RN6911 and RN6911-HLA), respectively. The survival rate of mice from different groups was assessed at the indicated time points. It was shown that restoration of α -toxin could partially recover lethality (Fig. 3). These results suggested that α -toxin played an important role in the intracranial infection causing abscesses.

Discussion

Intracranial abscesses caused by bacteria are usually associated with high mortality rate. *S. aureus* is a common cause of various kinds of infections, including intracranial abscesses. However, there is no report about animal model of intracranial abscesses caused by *S. aureus*. Moreover, the effect of a known regulator in *S. aureus* in intracranial infection has not been determined. In our study, we constructed a murine model of intracranial abscesses caused by *S. aureus*. We also demonstrated that agr, the quorum sensing system, in *S. aureus* was essential for causing intracranial infection. Furthermore, our results revealed that α -toxin regulated by agr system played an important role in the process of intracranial abscesses.

There are some constructed *S. aureus* infection models including arthritis, pneumonia, acute peritonitis, endocarditis, and skin infection.¹⁷⁻²⁶ Until now, there are few reports about intracranial infection model caused by *S. aureus*. In this study, we constructed a murine model by injecting *S. aureus* 8325-4 into the frontal cortex. Mice challenged with *S. aureus* showed some clinical signs of illness. The number of white blood cells and the rectal temperature were not significantly altered after injecting *S. aureus* suggesting that the sign of murine intracranial abscess was not exactly the same as human intracranial abscess. This may be due to difference in resistance to bacteria between mice and humans. However, the percentage of neutrophil in mice challenged with *S. aureus* was significantly higher than that of control group. This result is in accordance with the clinical sign seen in patients.

Agr system is well studied in *S. aureus* and is considered essential for infection.^{10,16,27} In line with expectations, we found that the mortality associated with agr-null strain was significantly decreased when compared to the wild type on intracranial abscesses model. Moreover, we also found that deletion of RNAIII suppressed virulence of *S. aureus* infection.

S. aureus can express various kinds of exotoxins which are necessary for causing infection. α -Toxin is the major exotoxin of *S. aureus*, regulated by RNAIII.^{10,16,28,29} Our results showed that restoration of the level of α -toxin in agr-null strain partially recovered the mortality associated with *S. aureus* on the intracranial abscesses model. This result suggested that α -toxin should play an important role in intracranial infection. Moreover, some other molecules regulated by the agr system might also be involved in intracranial infection. The role of α -toxin and other molecules expressed by *S. aureus* in intracranial infection should be further investigated in the future.

Our study presents a murine intracranial abscesses model that can be used to investigate *S. aureus* infection in brain tissue. Further investigation on the role of agr system will increase our understanding of the intracranial abscesses caused by *S. aureus*.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

This research was supported by National Natural Science Funding (31271119, 81072074), Beijing Educational Committee Key Funding (KZ201110025033) and Beijing Organization Department of Municipal Committee Talent Funding (3034000004).

REFERENCES

1. Saito N, Aoki K, Sakurai T, et al. Linezolid treatment for intracranial abscesses caused by methicillin-resistant *Staphylococcus aureus*. Neurol Med Chir. 2010;50:515-7.
2. Kono Y, Prevedello DM, Snyderman CH, et al. One thousand endoscopic skull base surgical procedures demystifying the infection potential: incidence and description of

- postoperative meningitis and brain abscesses. *Infect Control Hosp Epidemiol.* 2011;32:77-83.
3. Dashti SR, Baharvahdat H, Spetzler RF, et al. Operative intracranial infection following craniotomy. *Neurosurg Focus.* 2008;24:E10.
 4. Fulkerson DH, Sivaganesan A, Hill JD, et al. Progression of cerebrospinal fluid cell count and differential over a treatment course of shunt infection. *J Neurosurg Pediatr.* 2011;8:613-9.
 5. Mathisen GE, Johnson JP. Brain abscess. *Clin Infect Dis.* 1997;25:763-79.
 6. Kim JH, Desai NS, Ricci J, et al. Factors contributing to ventriculostomy infection. *World Neurosurg.* 2012;77: 135-40.
 7. Sifri CD, Park J, Helm GA, Stemper ME, Shukla SK. Fatal brain abscess due to community-associated methicillin-resistant *Staphylococcus aureus* strain USA300. *Clin Infect Dis.* 2007;45:e113-7.
 8. Wang KW, Chang WN, Shih TY, et al. Infection of cerebrospinal fluid shunts: causative pathogens, clinical features, and outcomes. *Jpn J Infect Dis.* 2004;57:44-8.
 9. Thoendel M, Kavanaugh JS, Flack CE, Horswill AR. Peptide signaling in the staphylococci. *Chem Rev.* 2010;111:117-51.
 10. Yarwood JM, Schlievert PM. Quorum sensing in *Staphylococcus* infections. *J Clin Invest.* 2003;112:1620-5.
 11. Inoshima I, Inoshima N, Wilke GA, et al. A *Staphylococcus aureus* pore-forming toxin subverts the activity of ADAM10 to cause lethal infection in mice. *Nat Med.* 2011;17:1310-4.
 12. LaFrance-Corey RG, Howe CL. Isolation of brain-infiltrating leukocytes. *J Vis Exp.* 2011;52.
 13. Liu GY, Essex A, Buchanan JT, et al. *Staphylococcus aureus* golden pigment impairs neutrophil killing and promotes virulence through its antioxidant activity. *J Exp Med.* 2005;202:209-15.
 14. Dinges MM, Orwin PM, Schlievert PM. Exotoxins of *Staphylococcus aureus*. *Clin Microbiol Rev.* 2000;13:16-34.
 15. Labandeira-Rey M, Couzon F, Boisset S, et al. *Staphylococcus aureus* Panton-Valentine leukocidin causes necrotizing pneumonia. *Science.* 2007;315:1130-3.
 16. Yarwood JM, Bartels DJ, Volper EM, Greenberg EP. Quorum sensing in *Staphylococcus aureus* biofilms. *J Bacteriol.* 2004;186:1838-50.
 17. Keffer J, Probert L, Cazlaris H, et al. Transgenic mice expressing human tumour necrosis factor: a predictive genetic model of arthritis. *EMBO J.* 1991;10:4025-31.
 18. Gong JH, Ratkay LG, Waterfield JD, Clark-Lewis I. An antagonist of monocyte chemoattractant protein 1 (MCP-1) inhibits arthritis in the MRL-lpr mouse model. *J Exp Med.* 1997;186:131-7.
 19. Chen J, Feng G, Guo Q, et al. Transcriptional events during the recovery from MRSA lung infection: a mouse pneumonia model. *PLoS ONE.* 2013;8:e70176.
 20. Ginsberg HS, Moldawer LL, Sehgal PB, et al. A mouse model for investigating the molecular pathogenesis of adenovirus pneumonia. *Proc Natl Acad Sci U S A.* 1991;88:1651-5.
 21. Leyboldt JK, Kamerath CD, Gilson JF. Acute peritonitis in a C57BL/6 mouse model of peritoneal dialysis. *Adv Perit Dial.* 2007;23:66-70.
 22. Takahashi K, Gordon J, Liu H, et al. Lack of mannose-binding lectin-A enhances survival in a mouse model of acute septic peritonitis. *Microbes Infect.* 2002;4:773-84.
 23. Xiong YQ, Willard J, Kadurugamuwa JL, Yu J, Francis KP, Bayer AS. Real-time *in vivo* bioluminescent imaging for evaluating the efficacy of antibiotics in a rat *Staphylococcus aureus* endocarditis model. *Antimicrob Agents Chemother.* 2005;49:380-7.
 24. Gibson GW, Kreuser SC, Riley JM, et al. Development of a mouse model of induced *Staphylococcus aureus* infective endocarditis. *Comp Med.* 2007;57:563-9.
 25. Kennedy AD, Bubeck Wardenburg J, Gardner DJ, et al. Targeting of alpha-hemolysin by active or passive immunization decreases severity of USA300 skin infection in a mouse model. *J Infect Dis.* 2010;202:1050-8.
 26. Brown EL, Dumitrescu O, Thomas D, et al. The Panton-Valentine leukocidin vaccine protects mice against lung and skin infections caused by *Staphylococcus aureus* USA300. *Clin Microbiol Infect.* 2009;15:156-64.
 27. Vuong C, Saenz HL, Götz F, Otto M. Impact of the agr quorum-sensing system on adherence to polystyrene in *Staphylococcus aureus*. *J Infect Dis.* 2000;182:1688-93.
 28. Queck SY, Jameson-Lee M, Villaruz AE, et al. RNAIII-independent target gene control by the agr quorum-sensing system: insight into the evolution of virulence regulation in *Staphylococcus aureus*. *Mol Cell.* 2008;32:150-8.
 29. Cheung AL, Bayer AS, Zhang G, Gresham H, Xiong YQ. Regulation of virulence determinants *in vitro* and *in vivo* in *Staphylococcus aureus*. *FEMS Immunol Med Microbiol.* 2004;40:1-9.
 30. Novick RP. Properties of a cryptic high-frequency transducing phage in *Staphylococcus aureus*. *Virology.* 1967;33:155-66.
 31. Peng HL, Novick RP, Kreiswirth B, Kornblum J, Schlievert P. Cloning, characterization, and sequencing of an accessory gene regulator (agr) in *Staphylococcus aureus*. *J Bacteriol.* 1988;170:4365-72.
 32. Liu Y, Mu C, Ying X, et al. RNAIII activates map expression by forming an RNA-RNA complex in *Staphylococcus aureus*. *FEBS Lett.* 2011;585:899-905.