



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Letter to the Editor

Frequency of CCR5 genotypes in HIV-infected patients in Roraima, Brazil



Dear Editor,

The expression of CCR5 gene, located on the short arm of chromosome three (3p21.31), leads to the formation of a protein that acts as a receptor for several chemokines.¹ A deletion of 32 base pairs gives rise to an important Rs333 polymorphism. The heterozygous genotype CCR5/ Δ 32 produces partially functional receptors while the homozygous recessive Δ 32/ Δ 32 features a non-functional receptor.²

Information about the prevalence of Rs333 is an essential aspect of medical decisions, mainly for CCR5 blockers prescription to HIV patients. Furthermore, immunogenetics studies that include the prevalence of CCR5 genes or other genetic factors in the population of Roraima State, Brazil, were not found in the literature.

To determine the prevalence of Rs333 polymorphism located in the CCR5 gene, we designed a cross-sectional study with HIV-infected patients living in Roraima, who had blood sent for analyses at the Central Laboratory of Roraima State (LACEN/RR). The Ethics Research Committee (REC) from the Federal University of Roraima (COEP/UFRR) approved this research (CAAE number: 15629013.8.0000.5302). Subsequently, 5 mL of blood samples were collected from 120 HIV-1 infected patients between January 2014 and February 2014. Genomic DNA was extracted using the automated system QIAcube, with QIAamp DNABlood Kit (QIAGEN, Germany) followed by PCR amplification as described previously³ and agarose gel electrophoresis. Three of these samples were rejected. Thus, our sampling represents more than 10% of all HIV-infected patients, including patients from 12 out of 14 municipalities in the Roraima state.

Two genotypes were found in this group of patients: 106 patients had CCR5/CCR5 represented by a single fragment of 137 bp, and 11 patients had CCR5/CCR5 Δ 32 characterized by two fragments, one of 137 bp and another of 105pb. Nevertheless, the CCR5 Δ 32/CCR5 Δ 32 genotype was not found in this study (Table 1).

Roraima is one of the newest states of the Brazilian Federation and one of the principal mining centers in the country during the 80s and 90s, which attracted Brazilians from other regions contributing to the population miscegenation.⁴ The Δ 32 allele has its origin in Europe; in Brazil it has different frequencies varying from 4.2% in the North region, to 6.4% in the South and 0.2% in Native americans.⁵ That frequency in the

Table 1 – Occurrence of genotypes, genotype frequency and allele frequency of the CCR5 gene in Roraima, Brazil.

Genetic profile	Frequency Patient (n = 117)
Genotype	
CCR5/CCR5	106 (90.6%)
CCR5/CCR5 Δ 32	11 (9.4%)
CCR5 Δ 32/CCR5 Δ 32	0
Genotypic frequency	
CCR5/CCR5	0.906
CCR5/CCR5 Δ 32	0.094
CCR5 Δ 32/CCR5 Δ 32	0
Allele frequency	
CCR5	0.954
CCR5 Δ 32	0.046

North region, where Roraima is located, is similar to what was found in this study (4.6%).

Due to the undoubted protective effect of CCR5 Δ 32 against HIV infection and aids progress, that project was not designed as an inference study to evaluate this marker as an immunogenetic protective factor in Roraima's HIV-infected patients. Thus, healthy controls were not included. Yet, to our knowledge, this is the first study about the prevalence of immunogenetic factors in HIV-infected patients from Roraima State.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

This work was supported by Instituto L eonidas e Maria Deane/FIOCRUZ – AM and LACEN-RR.

REFERENCES

- Liu R, Paxton WA, Choe S, et al. Homozygous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell*. 1996;86:367–77, [http://dx.doi.org/10.1016/S0092-8674\(00\)80110-5](http://dx.doi.org/10.1016/S0092-8674(00)80110-5).

2. Alkhatib G. The biology of CCR5 and CXCR4. *Curr Opin HIV AIDS*. 2009;4:96–103, <http://dx.doi.org/10.1097/COH.0b013e328324bbec>.
3. De Farias JD, Santos MG, De França AK, et al. Distribution of the CCR5delta32 allele (gene variant CCR5) in Rondônia, Western Amazonian region, Brazil. *Genet Mol Biol*. 2012;35:27–31, <http://dx.doi.org/10.1590/S1415-47572012005000003>.
4. Magno A, Diniz A. *Vertiginoso crescimento populacional de. Cad Geogr*. 2005;15:23–44.
5. Vargas AE, Marrero AR, Salzano FM, Bortolini MC, Chies JAB. Frequency of CCR5delta32 in Brazilian populations. *Braz J Med Biol Res*. 2006;39:321–5, <http://dx.doi.org/10.1590/S0100-879X2006000300002>.

André de Lima Guerra Corado^{a,*},
George Allan Villarouco da Silva^a,
Renato Augusto Carvalho Leão^b, Fabiana Granja^c,
Felipe Gomes Naveca^a

^a Instituto Lêonidas e Maria Deane/Fiocruz Amazonas, Manaus, AM, Brazil

^b Laboratório Central de Saúde Pública de Roraima (LACEN/RR), Boa Vista, RR, Brazil

^c Laboratório de Biologia Molecular, Universidade Federal de Roraima (UFRR), Boa Vista, RR, Brazil

* Corresponding author.

E-mail address: andrecorado@gmail.com (A.d.L.G. Corado).

Received 25 November 2015

Accepted 5 January 2016

Available online 2 March 2016

1413-8670/© 2016 Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)
<http://dx.doi.org/10.1016/j.bjid.2016.01.001>