



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

***Bartonella* species pathogenic for humans infect pets, free-ranging wild mammals and their ectoparasites in the Caatinga biome, Northeastern Brazil: a serological and molecular study**



Mariana Campos Fontalvo^a, Alexsandra Rodrigues de Mendonça Favacho^b,
Andreina de Carvalho Araujo^a, Naylla Mayana dos Santos^a,
Glauber Meneses Barboza de Oliveira^a, Daniel Moura Aguiar^c,
Elba Regina Sampaio de Lemos^b, Mauricio Claudio Horta^{a,*}

^a Universidade Federal do Vale do São Francisco (Univasf), Laboratório de Doenças Parasitárias, Petrolina, PE, Brazil

^b Instituto Oswaldo Cruz (FIOCRUZ), Laboratório de Hantavírus e Rickettsioses, Rio de Janeiro, RJ, Brazil

^c Universidade Federal de Mato Grosso, Laboratório de Virologia e Rickettsioses, Cuiabá, MT, Brazil

ARTICLE INFO

Article history:

Received 4 October 2016

Accepted 3 February 2017

Available online 27 February 2017

Keywords:

Bartonella

Flea

Tick

Lice

Brazil

ABSTRACT

This study verified the occurrence of *Bartonella* spp. in dogs, cats, wild mammals and their ectoparasites in Petrolina and Lagoa Grande Counties, Pernambuco, located in a semi-arid region in Northeastern Brazil. Anti-*Bartonella* spp. antibodies were detected by indirect immunofluorescence assay (IFA) in 24.8% of dogs (27/109) and in 15% of cats (6/40). *Bartonella* sp. DNA was identified by PCR performed on DNA extracted from blood and ectoparasites using primers targeting *Bartonella* sp. *gltA* and *ribC* genes in 100% (9/9) of *Pulex irritans* from *Cerdocyon thous*, 57.4% (35/61) of *P. irritans* from dogs, 2.3% (1/43) of *Ctenocephalides felis felis* from dogs, 53.3% (24/45) of *C. felis felis* from cats, and 10% (1/10) of *Polyplax* spp. from *Thrichomys apereoides*. DNA sequencing identified *Bartonella clarridgeiae* and *Bartonella henselae* in *C. felis felis* from cats, *Bartonella rochalimae* in *P. irritans* from dog and *C. thous*, and *Bartonella vinsoni berkhofii* in *P. irritans* from dog.

© 2017 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: horta.mc@hotmail.com (M.C. Horta).

<http://dx.doi.org/10.1016/j.bjid.2017.02.002>

1413-8670/© 2017 Sociedade Brasileira de Infectologia. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Bartonella spp. comprises a group of emerging pathogens that are prevalent in a large variety of vertebrates and cause diversified symptoms and clinical manifestations.¹ Transmission predominantly occurs via a vector, and fleas, hematophagous lice, and ticks have confirmed vector competence.² *Bartonella* DNA has also been isolated from mites.³

Bartonella henselae is the main causal agent of bartonellosis in humans. Although cats are a major reservoir for human infections with these *Bartonella* species, this agent can infect many mammalian hosts. Thus, rodents and other wild mammals can act as natural reservoirs for numerous species of these bacteria and the presence of the bacteria have been described in these animals on five continents.^{3–7}

In Brazil, the circulation of the agent has been demonstrated in humans, cats, dogs, and wild animals using serological and molecular techniques. However, little information is available about the occurrence of *Bartonella* in this country, and to date no study has verified the occurrence of *Bartonella* spp. in the Caatinga biome. Therefore, in the present study we investigated the circulation of *Bartonella* spp. in domestic mammals (dogs and cats), free-ranging small wild mammals (rodents, marsupials, and canids) and their ectoparasites (fleas, ticks and hematophagous lice) in the Caatinga biome.

Materials and methods

Study site

This study was conducted in the Petrolina (9°19'41" S, 40°33'30" W) and Lagoa Grande (8°40'1" S, 40°8'42" W) municipalities, State of Pernambuco, Brazil (Fig. 1). This region has semi-arid tropical weather and is part of the Caatinga biome in Northeastern Brazil.⁸ The municipalities are located in the region of the São Francisco Valley and represent two distinct areas within a degraded environment.

Sample collection

A total of 77 small mammals were trapped in eight trials from August 2014 to May 2015. Each trial was in a different area, with four in each municipality and two in each season of the year (spring, summer, autumn, and winter). Animals were chemically immobilized as recommended by Mares-Guia.⁹ Blood samples were collected from these animals via the caudal vein or intra-cardiac puncture. Additionally, ectoparasites were collected from the animals and stored in 1.5 mL microtubes containing absolute ethyl alcohol (C₂H₅OH) at –20 °C prior to laboratory analysis. After collection of biological materials, the animals were marked by cutting the hair from the sacral region and were set free in the same point of capture after full recovery of consciousness.

Domicile dogs and cats were analyzed in rural dwellings around each collection area. Information including age, gender, county of origin, tick, flea and/or louse presence, history of ectoparasitism, and access to the forest was obtained. Blood samples were collected from the cephalic vein in tubes

containing EDTA, properly identified, and centrifuged at 3000 rpm for 15 min. Plasma and whole blood were stored in 1.5 mL microtubes at –20 °C prior to analysis.

Ticks, fleas, and lice were collected and conserved in absolute ethanol and stored at room temperature until identification, according to Linardi and Guimaraes,¹⁰ Barros-Battesti, Arzua and Bechara,¹¹ and Pereira et al.,¹² and molecular testing.

All procedures followed the ethical standards of animal experimentation established by the Committee on Ethics and Studies and Research at the Federal University of São Francisco Valley – CEDEP/Univasf (protocol number 9/021014) and by the Brazilian Institute of Environment and Renewable Natural Resources - IBAMA (protocol number 45764-1) according to the recommendations and laws regarding the maintenance of animal welfare.

Indirect immunofluorescence assay (IFA)

Plasma samples from dogs and cats were subjected to indirect immunofluorescent antibody assay (IFA) (Bion, IL, USA) to detect anti-*Bartonella* sp. antibodies as recommended by the manufacturer. A cutoff value of 1:64 was used.¹³

Polymerase chain reaction (PCR)

DNA was extracted from whole blood of dogs, cats, and wild animals and from individual ticks using a commercial kit (Promega, Madison, WI, USA) as recommended by the manufacturer. A negative DNA extraction control consisting of 100 µL of sterile distilled water was included in each batch of samples. Whole fleas and lice were individually subjected to DNA extraction by boiling at 100 °C for 20 min.¹⁴

All samples were individually processed by PCR using the primers Bhcs.781p (5'-GGG GAC CAG CTC ATG GTG G-3') and Bhcs.1137n (5'-AAT GCA AAA AGA ACA GTA AAC A-3'), which amplified a 380-bp fragment of the citrate synthase gene (*gltA*) of *Bartonella* spp., and BARTON-1 (5'-TAA CCG ATA TTG GTT GTG TTG AAG-3') and BARTON-2 (5'-TAA AGC TAG AAA GTC TGG CAA CAT AAC G-3'), which amplified a 580-bp fragment of the riboflavin synthase C gene (*ribC*). The PCR reactions were performed as described previously.^{15,16} The PCR products were stained with ethidium bromide and visualized by electrophoresis in a 1.5% agarose gel.

PCR products of the expected amplicon size were purified, and their forward and reverse nucleotide sequences were subjected to direct sequencing using the ABI Prism BigDye Terminator v.3.1 cycle sequencing kit (Applied Biosystems, CA, USA). Partial sequences obtained from this study were submitted to BLAST (Basic Local Alignment Search Tool) analysis to determine similarities to other *Bartonella* species sequences available in the GenBank database.¹⁷

Statistical analysis

Categorical variables are described as proportions with the respective 95% confidence interval, and compared by the Chi-square test (X^2) using the software Epi Info 7.1.




Fig. 1 – Location of the municipalities of Lagoa Grande and Petrolina, Pernambuco State, Brazil.

Results

Seventy-seven small wild mammals were trapped over 2944 trap nights, showing a success rate of 2.6%. The mammals were identified as Rodentia [*Thrichomys apereoides* (30 specimens), *Wiedomys pyrrhorhinus* (3), *Galea spixii* (3), and *Calomys expulsus* (2)], Marsupialia [*Didelphis albiventris* (27), *Monodelphis domestica* (10), and *Gracilinanus agilis* (1)], and Canid [*Cercocyon thous* (1)]. By season, 15 mammals were trapped in winter (September 2014), 17 in spring (November 2014), 20 in summer (January 2015), and 25 in autumn (May 2015).

Hematophagous ectoparasites were found in 51.94% (40/77) of the wild animals. Among Rodentia, 52.63% (20/38) of the animals were infested: *T. apereoides* by *Amblyomma* sp. (larvae), *Amblyomma auricularium* (nymphs), Argasidae (larvae), and *Polyplax* sp. (adult) and *W. pyrrhorhinus* by *Amblyomma* sp. (larvae). Among Marsupialia, 50% (19/38) of the animals were infested: *D. albiventris* by *Amblyomma* sp. (larvae), *A. auricularium* (nymphs), *Amblyomma dubitatum* (nymphs), Argasidae (larvae), *Ctenocephalides felis felis* (adults), *M. domestica* by *Amblyomma* sp. (larvae), *A. auricularium* (nymphs), and Argasidae (larvae). Among Canidae, 100% (1/1) of the animals were infested: *C. thous* by *Pulex irritans* (adults). There was no infestation in *C. expulsus*, *G. spixii*, and *G. agilis*.

Approximately 56% of the dogs (61/109) were infested with *Rhipicephalus sanguineus* adult ticks and 19.26% (21/109) with *C. felis felis* (11/109) and/or *P. irritans* (16/109) fleas. Fifteen dogs (13.76%) were parasitized by both ticks and fleas. Around 22% of the cats (9/40) were infested with *C. felis felis* fleas.

Anti-*Bartonella* sp. antibodies were detected in 24.77% (27/109) of the dogs and 15% (6/40) of the cats. No significant differences were found among any of the variables analyzed ($p > 0.05$).

Bartonella sp. DNA was identified in nine (100%) *P. irritans* fleas from *C. thous*, one (10%) *Polyplax* sp. louse from *T. apereoides*, one (2.32%) *C. felis felis* and 35 (57.37%) *P. irritans* from dogs, and 24 (53.33%) *C. felis felis* from cats. All ticks tested negative by PCR (Table 1).

One dog was infested simultaneously with positive *C. felis felis* and *P. irritans*. Two dogs whose fleas were positive were reactive to the IFA. Fleas from eight (88.88%) cats showed *Bartonella* sp. DNA, of which two animals had titers of antibodies detectable by the IFA.

Partial sequences of the *gltA* gene were shown by the BLAST analysis to be 100% identical to the corresponding sequences of *Bartonella clarridgeiae* (KJ170236), *Bartonella henselae* (HG965802), *Bartonella rochalimae* (DQ676484.1) and *Bartonella vinsoni berkhoffii* (AF143445.1) (Table 1).

Discussion

The study of wildlife allows for an improved understanding of aspects related to the maintenance of various etiological agents in the epidemiological chain of different diseases as well as their sustainability and viability in nature. In this respect, it is important to emphasize studies related to zoonotic diseases due to the massive anthropization suffered

Table 1 – *Bartonella* spp. infection in hematophagous ectoparasites in wild and domestic mammals in a semi-arid region of Pernambuco, Brazil.

Host	% infestation (presence of ectoparasite/total)	Ectoparasites	Infection of <i>Bartonella</i> spp. by PCR	DNA sequencing
<i>Thrichomys apereoides</i>	63.3 (19/30)	<i>Amblyomma</i> sp.	0/3	–
		<i>A. auricularium</i>	0/23	–
		Argasidae	0/8	–
		<i>Polyplax</i> sp.	1/20	no
<i>Wiedomys pyrrhorhinus</i>	33.3 (1/3)	<i>Amblyomma</i> sp.	0/1	–
<i>Galea spixii</i>	0 (0/3)	–	0/0	–
<i>Calomys expulsus</i>	0 (0/2)	–	0/0	–
<i>Didelphis albiventris</i>	(15/27)	<i>Amblyomma</i> sp.	0/13	–
		<i>A. auricularium</i>	0/1	–
		<i>A. dubitatum</i>	0/4	–
		Argasidae	0/5	–
		<i>C. felis felis</i>	0/3	–
<i>Monodelphis domestica</i>	40.0 (4/10)	<i>Amblyomma</i> sp.	0/2	–
		<i>A. auricularium</i>	0/1	–
		Argasidae	0/1	–
<i>Gracilinanus agilis</i>	0 (0/1)	–	0/0	–
<i>Cerdocyon thous</i>	100 (1/1)	<i>P. irritans</i>	9/9	<i>B. rochalimae</i>
<i>Canis familiaris</i>	61.5 (67/109)	<i>R. sanguineus</i>	0/230	–
		<i>C. felis felis</i>	1/43	no
		<i>P. irritans</i>	35/61	<i>B. rochalimae</i> , <i>B. vinsoni berkhofii</i>
		<i>C. felis felis</i>	24/45	<i>B. clarridgeae</i> , <i>B. henselae</i>
<i>Felis catus</i>	22.5 (9/40)	<i>C. felis felis</i>	24/45	

by originally sylvan areas.¹⁸ In this study, a data survey of fauna ectoparasites was of great relevance due to the vector transmission of *Bartonella* spp.¹⁹

The capture success rate in this study (2.6%) was similar to that obtained by Dantas-Torres et al.²⁰ (1.7%) in the Atlantic Forest fragments in the State of Pernambuco, which is a biome that has suffered severe degradation similar to the Caatinga. However, the success rate was lower than that achieved by Guimarães²¹ (6.1%) in Serra das Confusões National Park, State of Piauí, which is an environmental preservation area with low human intervention rates. In line with these figures, Fonseca²² compared a non-degraded area to another degraded area in the State of Minas Gerais and obtained success rates of 6.2% and 2.9%, respectively.

Eight different mammalian species were detected, which was equivalent to approximately 5.4% of the total number of naturally occurring mammalian species in the Caatinga ($n=148$).²³ The most frequently caught species was *T. apereoides*, representing 39% (30/77) of the wild animals in the present study. This species was equivalent to 79% (30/38) of the rodents, which was similar to the 80% reported in Serra das Confusões National Park.²¹

We obtained a small number of animals infested with fleas in the present study compared to similar studies in both domestic and wild animals.^{24,25} This result may be related to climatic factors of the study site, which is located in the Drought Polygon region that presents features such as a negative water balance (resulting from an annual rainfall less than 500 mm), average insolation of 2800 h/year, annual average temperature of approximately 28° C (reaching up to 44.1° C in summer and 39.1° C in winter) and average relative humidity of 50%.^{26,27} These factors indicate that the site of the present study is a relatively hostile environment for order Siphonaptera, which generally prefers a temperature of

28 ± 1° C and a relative humidity of approximately 75% for its development.²⁸

Although no positive diagnosis was obtained in wild animals blood samples, it was possible to demonstrate the wild cycle of *Bartonella* spp. in the Caatinga by the presence of bacterial DNA in the fleas and blood-sucking lice that parasitized *C. thous* and *T. apereoides*, respectively.

The presence of *Bartonella* spp. in *Polyplax* spp. has been previously described in Taiwan.²⁴ The same study also detected DNA of the agent in fleas from wild animals but not in their ticks, in line with the results in our study. More recently, *Bartonella vinsonii* subsp. *arupensis* was identified in different species of wild rodents collected in Mato Grosso do Sul, which is the Brazilian state most covered by the Pantanal biome.²⁹

None of the dogs participating in the present study had any apparent sign of disease. Among the seropositive animals (titers ≥ 64), two had a positive diagnosis of *Bartonella* sp. DNA in their fleas but not in their blood samples. Dogs act as accidental reservoirs of some species of *Bartonella*. However, asymptomatic dogs rarely show persistent bacteremia.^{30,31}

The present study confirmed the occurrence of *Bartonella* spp. by detecting antibodies in dogs and cats and the presence of bacterial DNA in ectoparasites collected from domestic and wild animals. These results suggested that the hosts may harbor the microorganism, although *Bartonella* DNA was not found in any blood samples. Thus, the presence or absence of *Bartonella* spp. in the host does not necessarily reflect the condition of their fleas.^{32,33} Negative PCR results in the blood are often indicative of the bacterial DNA concentration in the sample, which may be insufficient for detection by the technique.^{34,35} Additionally, PCR of blood samples is commonly not successful due to the inhibitory factor hemin.³⁶ Furthermore, *Bartonella* causes a characteristic, cyclic bacteremia.³⁷

All positive samples tested by PCR targeting the *ribC* gene were also positive by PCR targeting the *gltA* gene, but not vice versa. This discrepancy may be due to variability in the sensitivity of the assay depending on the primer used or the primer targeting *gltA* may have been more comprehensive for the detection of a much wider range of *Bartonella* species that the primer targeting *ribC*.^{15,16,38}

The finding that an animal could be PCR-positive and IFA-negative may suggest that the infection is recent, the individual has not produced an IgG antibody level detectable by the test, the animal is immunosuppressed, or the infection is by a *Bartonella* species that does not present cross-reactivity with *B. henselae*.³⁹ This possibility may also explain why fleas from an animal are positive for *Bartonella* DNA when the animal does not have detectable antibodies by IFA as was observed in some animals in this study.

A total of 24.77% (34/109) of the dogs had a suggestive diagnosis for infection by *Bartonella* spp., and PCR-positive fleas were present on 42.3% (9/21) of the infested dogs. Among these, two dogs were both seropositive and had positive fleas. A lower prevalence was described by Brenner et al.⁴⁰ in stray dogs in São Paulo (9.3%, 11/118 animals). In Botucatu, state of São Paulo, only 3.5% of 198 dogs were seropositive for *Bartonella* spp.³⁰ In a study conducted in Peru, a 63% (68/108) seropositivity rate for *Bartonella* spp. was detected in asymptomatic dogs. The age group was shown to be a risk factor for animals aged less than one year,⁴¹ which was in contrast to the present study in which there was no such correlation. Of the 40 cats, six (15%) were positive in the IFA and seven (17.5%) in the PCR test of fleas. One cat was concurrently positive in the IFA and PCR test of fleas. Crissiuma et al.²⁵ reported the occurrence of *Bartonella* DNA in fleas from 20% (4/20) of infested cats, of which only one had detectable bacterial DNA in the blood.

The serological results in cats in this study are similar to those of Loureiro and Hagiwara,⁴² who detected 16% positivity for anti-*B. henselae* antibodies in 200 domicile cats in São Paulo. Among studies conducted with cats in Brazil, the seroreactivity ranged from 1.6% in domestic cats and cats from the Zoonosis Control Center of São Luís, Maranhão, and Cuiabá and Várzea Grande, Mato Grosso^{43,44} to up to 97% in cats from shelters in Vassouras, Rio de Janeiro,⁴⁵ using molecular and/or serological methods. The studies were performed in the States of São Paulo [4.3%,⁴⁶ 30%⁴⁷], Rio de Janeiro [42.5%,²⁵ 35.7%,¹³ and 56.1%⁴⁸], Mato Grosso [2.2%⁴⁹] and Rio Grande do Sul [25.5%⁵⁰].

Conclusions

The results of the present study confirm for the first time the occurrence of *Bartonella* in the Caatinga biome and in the semi-arid region and the identification of at least four different species (*B. clarridgeiae*, *B. henselae*, *B. rochalimae* and *B. vinsoni berkhofii*) in the same geographic region of Northeastern Brazil. This is the first report of *Bartonella* spp. in *Polyplax* sp. and *Pulex* sp. in Brazil. Further investigations are needed to identify the prevalent *Bartonella* species, to verify the vector competence of these flea species, and to elucidate their epidemiology at the study site.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank Ana Isabel A. Santos, Lais F. Santos, Josenilton R. Santos, Ivo W. G. da Silva, and Dália M. R. Machado for their valuable help during the field work and laboratory tests and the Centro de Conservação e Manejo de Fauna (Cema-Fauna) for confirmation of the identification of wild mammalian species. This work was supported by the Brazilian Research Funding Agency (CNPq).

REFERENCES

1. Boulouis HJ, Chang CC, Henn JB, Kasten RW, Chomel BB. Factors associated with the rapid emergence of zoonotic *Bartonella* infections. *Vet Res.* 2005;36:383-410.
2. Reis C, Cote M, Le Rhun D, Lecuelle B, Levin ML, Vayssier-Taussat M, et al. Vector competence of the tick *Ixodes ricinus* for transmission of *Bartonella birtlesii*. *PLoS Negl Trop Dis.* 2011;5:e1186.
3. Kim CM, Kim JY, Yi YH, Lee MJ, Cho MR, Shah DH, et al. Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *J Vet Sci.* 2005;6:327-34.
4. Birtles RJ, Harrison TG, Molyneux DH. *Grahamella* in small woodland mammals in the U.K.: isolation, prevalence and host specificity. *Ann Trop Med Parasitol.* 1994;88:317-27.
5. Yamamoto K, Chomel BB, Lowenstine LJ, Kikuchi Y, Phillips LG, Barr BC, et al. *Bartonella henselae* antibody prevalence in free-ranging and captive wild felids from California. *J Wildl Dis.* 1998;34:56-63.
6. Merhej V, Croce O, Robert C, Rolain J, Raoult D. Genome sequence of *Bartonella rattaustraliani*, a bacterium isolated from an Australian rat. *J Bacteriol.* 2012;194:7012.
7. Meheretu Y, Leirs H, Welegerima K, Breno M, Tomas Z, Kidane D, et al. *Bartonella* prevalence and genetic diversity in small mammals from Ethiopia. *Vector Borne Zoonotic Dis.* 2013;13:164-75.
8. IBGE (Instituto Brasileiro de Geografia e Estatística). Manual técnico da vegetação brasileira. Rio de Janeiro: Fundação IBGE; 2012. p. 272.
9. Mares-Guia MAMM, Moreira N, Monteiro RV, Barreira JD, Roque AL. Técnicas de Contenção Química de Pequenos Mamíferos Silvestres: princípios éticos e do bem-estar animal no campo. In: Lemos ERS, D'Andrea, editors. Trabalho de Campo com Animais: procedimentos, riscos e biossegurança. Rio de Janeiro: Editora Fiocruz; 2014. p. 95-104.
10. Linardi PM, Guimarães LR. Sifonápteros do Brasil, Museu de Zoologia. São Paulo: USP/FAPESP; 2000. p. 291.
11. Barros-Battesti DM, Arzua M, Bechara GH. Carrapatos de importância médico-veterinária da região neotropical: um guia ilustrado para identificação de espécies. São Paulo: Instituto Butantan; 2006. p. 223.
12. Pereira JS, Fonseca ZAAS, Gadelha ICN, Paiva KAR, Ahid SMM. Parasitismo por *Polyplax spinulosa* Burmeister, 1839 (Anoplura) em ratos Wistar, *Rattus norvegicus* Berkenhout, 1769. *Rev Bras Hig San An.* 2015;9:105-10.
13. Lamas CC, Mares-Guia MA, Rozental T, Moreira N, Favacho ARM, Barreira J, et al. *Bartonella* spp. infection in HIV positive individuals, their pets and ectoparasites in Rio de Janeiro,

- Brazil: serological and molecular study. *Acta Trop*. 2010;115:137–41.
14. Horta MC, Labruna MB, Pinter A, Linardi PM, Schumaker TTS. *Rickettsia* infection in five areas of the state of São Paulo, Brazil. *Mem Inst Oswaldo Cruz*. 2007;102:793–801.
 15. Norman AF, Regnery R, Jameson P, Greene C, Krause DC. Differentiation of *Bartonella*-like isolates at the species level by PCR-restriction fragment length polymorphism in the citrate synthase gene. *J Clin Microbiol*. 1995;33:1797–803.
 16. Johnson G, Ayers M, McClure SCC, Richardson SE, Tellier R. Detection and identification of *Bartonella* species pathogenic for humans by PCR amplification targeting the riboflavin synthase gene (*ribC*). *J Clin Microbiol*. 2003;41:1069–72.
 17. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215:403–10.
 18. Barbosa AD, Martins NRS, Magalhães DF. Zoonoses e saúde pública: riscos da proximidade humana com a fauna silvestre. *Cienc Vet Trop*. 2011;14:1–9.
 19. Chomel BB, Boulouis HJ. Zoonotic diseases caused by bacteria of the *Bartonella* genus: new reservoirs? New vectors? *Bull Acad Natl Med*. 2005;189:465–80.
 20. Dantas-Torres F, Aléssio FM, Siqueira DB, Mauffrey JF, Marvulo MFV, Martins TF, et al. Exposure of small mammals to ticks and rickettsiae in Atlantic Forest patches in the metropolitan area of Recife, Northeastern Brazil. *Parasitology*. 2012;139:83–91.
 21. Guimarães MF. Levantamento da Infecção por *Rickettsia* spp. e *Coxiella burnetii* em Mamíferos Silvestres e Pequenos Ruminantes na Região do Parque Nacional da Serra das Confusões, Piauí. Petrolina: Dissertação (Mestrado em Ciência Animal) - Universidade Federal do Vale do São Francisco; 2014. p. 91.
 22. Fonseca MT. A Estrutura da Comunidade de Pequenos Mamíferos em um Fragmento de Mata Atlântica e Monocultura de Eucalipto: A importância da matriz de habitat. Belo Horizonte: Dissertação (Mestrado em Ecologia, Conservação e Manejo de Vida Silvestre) - Universidade Federal de Minas Gerais; 1997. p. 60.
 23. Oliveira JA, Gonçalves PR, Bonvicino CR. Mamíferos da Caatinga. In: Leal IR, Tabarelli M, Silva JMC, editors. *Ecologia e Conservação da Caatinga*. Recife. 2003, 800 pp.
 24. Tsai YL, Chuang ST, Chang CC, Kass PH, Chomel BB. *Bartonella* species in small mammals and their ectoparasites in Taiwan. *Am J Trop Med Hyg*. 2010;83:917–23.
 25. Crissiuma A, Favacho A, Gershony L, Mendes-de-Almeida F, Gomes R, Mares-Guia A, et al. Prevalence of *Bartonella* species DNA and antibodies in cats (*Felis catus*) submitted to a spay/neuter program in Rio de Janeiro, Brazil. *J Feline Med Surg*. 2011;13:149–51.
 26. Brazil. Ministério da Integração Nacional. Relatório Final. Grupo de Trabalho Interministerial para redelimitação do Semi-árido Nordeste e do Polígono das Secas; 2005. Available from: http://www.mi.gov.br/c/document_library/get_file?uuid=090e3f78-bde3-4a1b-a46c-da4b1a0d78fa&groupId=10157
 27. INMET. Instituto Nacional de Meteorologia. Ministério da Agricultura, Pecuária e Abastecimento; 2015. Available from: http://www.inmet.gov.br/portal/arq/upload/BOLETIM-AGRO_MENSAL_201501.pdf
 28. Silva CS, Veronez VA, Castagnoli KC, Prette N, Borges FA, Miyasaka DS. Implantação de colônia de *Ctenocephalides felis felis* (Bouché, 1835) e determinação do período de desenvolvimento dos estágios imaturos sob condições controladas. *Ambiência: Rev Set Ciênc Agr Amb*. 2008;4:473–81.
 29. Favacho ARM, Andrade MN, Oliveira RC, Bonvicino CR, D'Andrea PS, de Lemos ERS. Zoonotic *Bartonella* species in wild rodents in the state of Mato Grosso do Sul, Brazil. *Microb Infect*. 2015;17:889–92.
 30. Diniz PPVP, Maggi RG, Schwartz DS, Cadenas MB, Bradley JM, Hegarty B, et al. Canine bartonellosis: serological and molecular prevalence in Brazil and evidence of co-infection with *Bartonella henselae* and *Bartonella vinsonii* subsp. *berkhoffii*. *Vet Res*. 2007;38:697–710.
 31. Breitschwerdt EB. Feline bartonellosis and cat scratch disease. *Vet Immunol Immunopathol*. 2008;123:167–71.
 32. Chomel BB, Kasten RW, Floyd-Hawkins K, Chi B, Yamamoto K, Roberts-Wilson J, et al. Experimental transmission of *Bartonella henselae* by the cat flea. *J Clin Microbiol*. 1996;34:1952–6.
 33. La Scola B, Davoust B, Boni M, Raoult D. Lack of correlation between *Bartonella* DNA detection within fleas, serological results, and results of blood culture in a *Bartonella*-infected stray cat population. *Clin Microbiol Infect*. 2002;8:345–51.
 34. Rubio AV, Ávila-Flores R, Osikowicz LM, Bai Y, Suzán G, Kosoy MY. Prevalence and genetic diversity of *Bartonella* strains in rodents from Northwestern Mexico. *Vector Borne Zoonotic Dis*. 2014;14:838–45.
 35. André MR, Dumler JS, Herrera HM, Gonçalves LR, Sousa KCM, Scorpio DG, et al. Assessment of a quantitative 5' nuclease real-time polymerase chain reaction using the nicotinamide adenine dinucleotide dehydrogenase gamma subunit (*nuoG*) for *Bartonella* species in domiciled and stray cats in Brazil. *J Feline Med Surg*. 2016;18:783–90.
 36. Akane A, Matsubara K, Nakamura H, Takahashi S, Kimura K. Identification of the heme compound co-purified with deoxyribonucleic acid (DNA) from bloodstains, a major inhibitor of polymerase chain reaction (PCR) amplification. *J Forensic Sci*. 1994;39:362–72.
 37. Harms A, Dehio C. Intruders below the radar: molecular pathogenesis of *Bartonella* spp. *Clin Microbiol Rev*. 2012;25:42–78.
 38. Buffet JP, Kosoy M, Vayssier-Taussat M. Natural history of *Bartonella*-infecting rodents in light of new knowledge on genomics, diversity and evolution. *Future Microbiol*. 2013;8:1117–28.
 39. Pretorius AM, Kelly PJ, Birtles RJ, Raoult D. Isolation of *Bartonella henselae* from a serologically negative cat in Bloemfontein, South Africa. *J S Afr Vet Assoc*. 1999;70:154–5.
 40. Brenner EE, Chomel BB, Singhasivanon OU, Namekata DY, Kasten RW, Kass PH, et al. *Bartonella* infection in urban and rural dogs from the tropics: Brazil, Colombia, Sri Lanka and Vietnam. *Epidemiol Infect*. 2013;141:54–61.
 41. Diniz PPPV, Morton BA, Tngrian M, Kachani M, Barrón EA, Gavidia CM, et al. Infection of domestic dogs in Peru by zoonotic *Bartonella* species: a cross-sectional prevalence study of 219 asymptomatic dogs. *PLoS Negl Trop Dis*. 2013;5:e2393.
 42. Loureiro VL, Hagiwara M. Levantamento de anticorpos anti-*Bartonella henselae* em felinos domiciliados na cidade de São Paulo, Estado de São Paulo e sua importância em saúde pública. *Rev Bras Ciênc Vet*. 2007;14:39–42.
 43. Braga MSCO, Diniz PPVP, André MR, Bortoli CP, Machado RZ. Molecular characterisation of *Bartonella* species in cats from São Luís, state of Maranhão, North-eastern Brazil. *Mem Inst Oswaldo Cruz*. 2012;107:772–7.
 44. Braga IA, Dias ISSO, Chitarra CS, Amuded AM, Aguiar DM. Molecular detection of *Bartonella clarridgeiae* in domestic cats from Midwest Brazil. *Braz J Infect Dis*. 2015;19:451–2.
 45. Souza AM, Almeida DNP, Guterres A, Gomes R, Favacho ARM, Moreira NS, et al. Bartonelose: análise molecular e sorológica em gatos do Rio de Janeiro – Brasil. *Rev Bras Ciênc Vet*. 2010;17:7–11.
 46. Bortoli CP, André MR, Seki MC, Pinto AA, Machado STZ, Machado RZ. Detection of hemoplasma and *Bartonella* species

- and co-infection with retroviruses in cats subjected to a spaying/neutering program in Jaboticabal, SP, Brazil. *Rev Bras Parasitol Vet.* 2012;21:219-23.
47. André MR, Denardia NCB, Sousa KCM, Gonçalves LR, Henrique PC, Ontivero CRGR, et al. Arthropod-borne pathogens circulating in free-roaming domestic cats in a zoo environment in Brazil. *Ticks Tick Borne Dis.* 2014;5: 545-51.
48. Kitada AAB, Favacho ARM, Oliveira RVC, Pessoa Junior AA, Gomes R, Honse CO, et al. Detection of serum antibodies against *Bartonella* species in cats with sporotrichosis from Rio de Janeiro, Brazil. *J Feline Med Surg.* 2014;16:308-11.
49. Miceli NG, Gavioli FA, Gonçalves LR, André MR, Sousa VR, Sousa KC, et al. Molecular detection of feline arthropod-borne pathogens in cats in Cuiabá, State of Mato Grosso, central-western region of Brazil. *Rev Bras Parasitol Vet.* 2013;22:385-90.
50. Staggemeier R, Venker CA, Klein DH, Petry M, Spilki FR, Cantarelli VV. Prevalence of *Bartonella henselae* and *Bartonella clarridgeiae* in cats in the south of Brazil: a molecular study. *Mem Inst Oswaldo Cruz.* 2010;200:873-8.