



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original article

Evaluation of the cervicovaginal environment in asymptomatic Human T-cell lymphotropic virus type 1 infected women



Alisson de Aquino Firmino^a, Adenilda Lima Lopes Martins^a, Luana Leandro Gois^{a,b},
Taiane Silva Paixão^a, Everton da Silva Batista^a, Bernardo Galvão-Castro^{a,b},
Maria Fernanda Rios Grassi^{id a,b,*}

^a Escola Bahiana de Medicina e Saúde Pública, Salvador, BA, Brazil

^b Fundação Oswaldo Cruz, Instituto Gonçalo Moniz, Salvador, BA, Brazil

ARTICLE INFO

Article history:

Received 14 September 2018

Accepted 7 February 2019

Available online 6 March 2019

Keywords:

HTLV-1

Inflammatory cytokines

Proviral load

Cervicovaginal cytopathology

ABSTRACT

Introduction: Human T-cell lymphotropic virus type 1 (HTLV-1) is sexually transmitted and causes persistent infection. This virus induces activation of the immune system and production of inflammatory cytokines. This study aimed to assess the cytokine profile and cytopathological findings in the cervicovaginal fluid of asymptomatic HTLV-1-infected women.

Methods: HTLV-1-infected and uninfected women were selected at the Centro de Atendimento ao Portador de HTLV in Salvador-Brazil. None of the included HTLV-1-infected women reported any HTLV-1-associated diseases. All volunteers underwent gynecological examination to collect cervicovaginal fluid. Cytokine quantification was performed using the Cytometric Bead Array (CBA) Human Th1/Th2/Th17 kit. Light microscopy was used to evaluate cervicovaginal cytopathology. In addition, proviral load in cervicovaginal fluid and peripheral blood was measured by real-time quantitative polymerase chain reaction.

Results: 112 women (63 HTLV-1-infected and 49 uninfected) were evaluated. No differences were found with respect to cytopathological cervicovaginal findings between the groups. IL-2, TNF, IL-4, IL-10, and IL-17 levels were significantly higher in cervicovaginal fluid of the HTLV-1-infected women than in uninfected women ($p < 0.05$). Conversely, IFN- γ was found to be lower in the HTLV-1-infected women ($p < 0.001$) compared to uninfected individuals. Cervicovaginal proviral load was detectable in 53% of the HTLV-1-infected women and was found to be consistently lower than the proviral load in peripheral blood.

Conclusions: HTLV-1 infection induces immune activation in cervicovaginal environment, characterized by elevated concentrations of Th1, Th2, and IL17 in the cervicovaginal fluid.

© 2019 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author.

E-mail address: fernanda.grassi@fiocruz.br (M.F. Grassi).

<https://doi.org/10.1016/j.bjid.2019.02.001>

1413-8670/© 2019 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Human T-cell lymphotropic virus type 1 (HTLV-1) is a retrovirus distributed worldwide, with endemic areas found in Africa, South and Central America, the Caribbean, Japan, Melanesia, and the Middle East. In Brazil, it is estimated that approximately 800,000 people are infected.¹ This virus is transmitted vertically from mother to child, mainly by breastfeeding, or horizontally through transfusion of blood, contaminated needles, or sexual intercourse. A recent study conducted in Salvador, Brazil, underscored the importance of the sexual route in HTLV-1 transmission.² In this city, HTLV-1-infection is more prevalent in women, reaching 10% of those over the age of 50.³

HTLV-1 is the etiological agent of adult T-cell leukemia/lymphoma (ATLL),⁴ HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP),⁵ HTLV-1 associated uveitis,⁶ and infective dermatitis in children.⁷ In addition, inflammatory diseases, such as keratoconjunctivitis sicca (KCS),⁸ bronchiectasis⁹ and arthritis¹⁰ are often associated with this viral infection.

Memory CD4⁺ T-lymphocytes are the main target cells for HTLV-1, although CD8⁺ T-lymphocytes, glial cells, and circulating dendritic cells may also represent targets for infection.¹¹⁻¹³ Immune system activation is a hallmark of HTLV-1 infection, resulting in spontaneous lymphoproliferation, increased expression of activation markers (HLA-DR, CD25), and exacerbated production of proinflammatory cytokines (IFN- γ , TNF) and chemokines (IL-8, CXCL9 and CXCL10).^{14,15} Studies indicate that elevated proviral load (>5% of infected cells) and immune activation are commonly found in the context of HTLV-1-associated diseases, as compared to asymptomatic HTLV-1 carriers.¹⁶⁻²⁰ However, asymptomatic individuals with low proviral load may also present pronounced immune activation.²¹

Few studies have evaluated the effect of HTLV-1 infection in the vaginal mucosa. Detection of HTLV-1 DNA in the cervical fluid of infected women was previously associated with cervicitis.²² In addition, presence of anti-HTLV-1 in the vaginal fluid has been described, including in asymptomatic HTLV-1 infected women.²³ This study evaluates the cervicovaginal environment of HTLV-1 infected women by assessing proviral load, cytopathological alterations, and cytokine levels.

Methods

Recruitment and study design

The present cross-sectional observational study was conducted at the Centro de Atendimento ao Portador de HTLV of the Escola Bahiana de Medicina e Saúde Pública, Salvador, Bahia, Brazil. Asymptomatic HTLV-1 infected women were sequentially included in the course of routine medical consultations from August 2014 to March 2016. All individuals were included if their neurologic examination was normal and no complaints for HTLV-1-associated diseases were reported. Uninfected women (controls) were selected from companions or relatives of patients who attended consultations. These

individuals were paired to HTLV-1-infected women matched for age, presence of comorbidities, smoking, contraceptive methods, and presence of other sexually transmitted infections. The sample size for the HTLV-1 asymptomatic group was calculated based on a 30% estimated prevalence of sexual dysfunction for HTLV-1 uninfected women, with an estimated prevalence ratio (PR) of 2.0 among both the HTLV-1-infected and uninfected women. Adopting an alpha error of 5% and power of 80%, the necessary sample size was determined to be 49 women in each group. The inclusion criteria consisted of age ranging from 20 to 50 years and report of sexual activity within four weeks preceding the consultation. Women with HTLV-1-associated diseases, those who were menopausal or diagnosed with depression, as well as those taking medication known to affect sexual desire (beta blockers, antidepressants, central nervous system depressants or anticholinergics) were excluded. HTLV-1 infection was diagnosed by Enzyme-Linked Immunosorbent Assay (ELISA) with Western Blot used for confirmation. This study was approved by the Institutional Research Board of the Escola Bahiana de Medicina e Saúde Pública (registered under protocol: CAAE 33098414.4.0000.5544) and all included women signed a term of informed consent.

Collection and analysis of samples

Following the routine medical consultation, demographic, medical, sexual, and gynecological data were obtained through specific standardized data collection forms, and physical and gynecological examinations were performed. Papanicolaou smears were collected from the ectocervix and endocervix using an Ayres spatula and cytobrush, respectively. Cotton swabs were used to collect vaginal fluid from the ectocervix, endocervix and vaginal walls for proviral load (PVL) measurement and cytokine quantification.

Cell abnormalities detected in the Papanicolaou smears were classified in accordance with the Bethesda System.²⁴ To measure PVL in the cervicovaginal fluid, swabs were placed in tubes containing 400 μ L of hydroxymethyl-ethylene diamine tetra acetic acid (Tris-EDTA) solution and stored at -20°C until use. For cytokine quantification, swabs were preserved in cryotubes containing 1 mL of sterile phosphate-buffered saline (PBS) stored at -70°C . Cytokine levels were assessed by flow cytometry using the Cytometric Bead Array (CBA) Human Th1/Th2/Th17 kit (Becton, Dickinson and Company, New Jersey, USA). Whole blood samples were collected in EDTA tubes and peripheral blood mononuclear cells (PBMC) were obtained by density gradient centrifugation and cryopreserved until use. HTLV-1 PVL in cervicovaginal cells and PBMCs was determined by real-time TaqMan PCR, as described elsewhere.²⁵

Statistical analyses

Data were expressed as medians and percentiles (25th and 75th) or means and standard deviation. Differences between HTLV-1 infected and uninfected women concerning family income, schooling, length of relationship, number of partners, parity, proviral load, and cytokine levels were assessed by the non-parametric Mann-Whitney *U* test, and differences in age were evaluated by the Student's *t*-test.

Table 1 – Sociodemographic profile of HTLV-1-infected and uninfected women.

Variable	HTLV-1-Infected (n = 63)	HTLV-1 uninfected (n = 49)	p-value
Age (years) ^a	34.68 (7.18)	35.61 (7.87)	0.52
Family income (no. of minimum wages) ^b	1.2 (1.0–2.0)	2.0 (1.0–2.25)	0.01
Educational level (years) ^b	10.0 (5.0–12.0)	12.0 (8.0–12.0)	0.08
Skin color n (%) ^c			0.14
Black	33 (52.4)	25 (51.0)	
Mixed	21 (33.3)	22 (44.9)	
White	9 (14.3)	2 (4.1)	
Marital status n (%) ^c			0.13
Married/Stable union	50 (79.4)	35 (71.4)	
Single	13 (20.6)	11 (22.4)	
Divorced/Separated	–	3 (6.1)	
Length of relationship (years) ^b	8.0 (3.75–12.0)	3.0 (1.15–7.5)	0.01
Number of partners ^b	4.0 (3.0–10.0)	4.0 (2.0–7.0)	0.38
Parity ^b	2.0 (1.0–3.0)	2.0 (0.5–2.0)	0.40

^a Data presented as mean (standard deviation) – p-value: Student's t test.

^b Data presented as median and interquartile range (p25–p75) – p-value: Mann–Whitney U test.

^c Data presented as frequency/proportion – p-value: Chi-square test.

Differences in qualitative variables (skin color, marital status and cytopathological/vaginal microbiota) were assessed using the Chi-square test. Spearman's rank correlation coefficient was used to determine associations between cytokine levels and proviral load. p-values less than 0.05 were considered statistically significant. All analyses were performed using GraphPad software version 5.0 and SPSS software version 17.0 for Windows.

Results

A total of 112 women (63 infected with HTLV-1 and 49 uninfected) were evaluated (Table 1). No significant differences between the groups were seen regarding sociodemographic

profile, number of partners, or parity. HTLV-1-infected women had longer relationships than uninfected women ($p = 0.01$) and the median income of HTLV-1-infected women (1.2 minimum wage) was lower compared to that of uninfected women (2 minimum wages) ($p = 0.01$). HTLV-1 proviral load was detected in 94% of the PBMC samples from the infected women, with a median of 28,665 copies/ 10^6 cells (IQR 4868–69,408 copies/ 10^6 cells).

Regarding cytopathological findings were similar among HTLV-1 infected or uninfected women with negative results for neoplasia ($p = 0.71$) (Table 2). Atypical squamous cells of undetermined significance (ASC-US) was identified in 1.6% and 2% of infected and uninfected women, respectively ($p = 0.86$). In addition, similar vaginal microbiota (*Lactobacillus vaginalis*, *Gardnerella vaginalis*/*Mobiluncus spp.*,

Table 2 – Frequency of cervicovaginal cytopathologic findings and HTLV-1 proviral load in cervicovaginal and PBMC samples.

Variable	HTLV-1-infected (n = 63)	HTLV-1 uninfected (n = 49)	p-value
Cervicovaginal cytopathology n (%) ^a			
Negative for neoplasia	61 (96.8)	48 (98.0)	0.71
ASC-US ^b	1 (1.6)	1 (2)	0.86
Unsatisfactory	1 (1.6)	0	0.38
Vaginal microbiota n (%) ^a			
<i>Lactobacillus vaginalis</i>	11 (17.5)	8 (16.3)	0.87
<i>Gardnerella vaginalis</i> / <i>Mobiluncus spp.</i>	13 (20.6)	15 (30.6)	0.23
Coccus/Bacillus	23 (36.5)	16 (32.7)	0.67
<i>Candida spp.</i>	16 (25.4)	9 (18.4)	0.37
<i>Trichomonas vaginalis</i>	–	1 (2)	0.25
HTLV-1 PVL in cervicovaginal fluid ^{c,d}	62 (0–2057)	NA	NA
HTLV-1 PVL in PBMC ^{c,e}	28,665 (4868–69,408)	NA	NA

^a Data presented as frequency/proportion – p-value: Chi-square test.

^b Atypical Squamous Cells of Undetermined Significance.

^c Data presented as median and interquartile range (p25–p75) – p-value: Mann–Whitney U test.

^d Considering 57 women, number of HTLV-1 copies/ 10^6 cells.

^e Considering 51 women, number of HTLV-1 copies/ 10^6 cells. PVL, proviral load; PBMC, peripheral blood mononuclear cells; NA, not applicable.

Coccus/Bacillus, *Candida spp.*, and *Trichomonas vaginalis*) were found in the two groups. HTLV-1 proviral load was detectable in 53% of the cervicovaginal samples from HTLV-1-infected women. The median cervicovaginal proviral load [62 copies/ 10^6 cells (IQR 0–2057 copies/ 10^6 cells)] was consistently lower than that in peripheral blood [28,665 copies/ 10^6 cells (IQR 4868–69,408)].

HTLV-1-infected women had significantly higher concentrations of IL-2, TNF, IL-10, IL-4, and IL-17 in the cervicovaginal fluid than uninfected women ($p < 0.05$). Conversely, these women presented lower concentrations of IFN- γ in the vaginal fluid compared to those uninfected by HTLV-1 ($p < 0.001$) (Fig. 1). The IFN- γ /IL-10 ratio in HTLV-1-infected women (8.87, IQR 8.77–9.10) was significantly lower than in uninfected




Fig. 1 – Levels of IL-2 (A), TNF (B), IFN- γ (C), IL-4 (D), IL-6 (E), IL-10 (F) and IL-17 (G) in cervicovaginal fluid of HTLV-1-infected ($n = 63$) and uninfected women ($n = 49$).

women (9.05, IQR 8.88–9.23), ($p < 0.001$). Cytokine levels were not found to be correlated with HTLV-1 proviral load in either the cervicovaginal fluid or in peripheral blood mononuclear cells.

Discussion

The present results indicate that, in addition to elevated levels of regulatory IL-10, both Th1 (IL-2 and TNF) and Th2 (IL-4) cytokines, as well as IL-17, were all higher in the cervicovaginal fluid of HTLV-1-infected women as compared to uninfected individuals. This immune activation of vaginal environment may be consequent to the presence of infected cells in the vaginal mucosa. Indeed, HTLV-1 proviral load was found to be detectable in the vaginal fluid of more than 50% of the infected women. Corroborating our results, Belec et al. found viral DNA in three out of 15 HTLV-1-infected women and suggested that the virus induced a local immune response that resulted in increased levels of HTLV-1 antibodies in the vaginal fluid.²³ In addition, the presence of HTLV DNA in cervical samples obtained from HTLV-1-infected sex workers was associated with the diagnosis of cervicitis.²²

It has been well established that HTLV-1 induces activation of the immune system, which is reflected by spontaneous proliferation of peripheral blood mononuclear cells and production of cytokines. Individuals with a diagnosis of HAM/TSP commonly present higher plasma levels of proinflammatory cytokines (e.g. IFN- γ , TNF, IL-6) and chemokines (CXCL9, CXCL10) than asymptomatic or uninfected individuals.^{14,16,26} Moreover, other HTLV-1-associated conditions, such as neurogenic bladder,²⁷ infective dermatitis¹⁷ and sicca syndrome,¹⁸ have also been reported in association with higher plasma levels of IFN- γ and IL-6.¹⁶ In addition, high production of IFN- γ is found in asymptomatic HTLV-1-infected individuals compared to uninfected controls.¹⁴ In the present study, low levels of IFN- γ and lower IFN- γ /IL-10 ratios were found in the cervicovaginal samples from the HTLV-1-infected group. The regulatory cytokine IL-10 has been reported to have an antagonist effect on the production of IFN- γ .²⁸ The predominance of IL-10 and TGF- β is commonly found in the mucosa of healthy individuals, which creates a regulatory milieu that maintains an immunological tolerance against antigens from microbiota and the external environment.^{29,30} The significantly higher IL-10 levels found in the cervicovaginal fluid of HTLV-1-infected women compared to uninfected women seems to suggest that the presence of the virus induced increased immune regulation.

Similarly, women infected with HIV, another retrovirus, also present increased levels of Th1 and Th2 cytokines in the vaginal fluid as compared to uninfected controls.^{30,31} Regarding IL-17, a study conducted in HIV-1-infected women found higher IL-17 concentrations in the vaginal fluid of women with sexually transmitted bacterial infections than in those without these types of infections, whereas women with *Candida* spp. had lower IL-17 concentrations compared to those without candidal infections.³²

It has been demonstrated that local immune activation induced by HIV in the vaginal environment may modulate

virus shedding in cervicovaginal secretions.³³ It is possible that HTLV-1 induces a similar immune activation, thereby increasing cytokine production *in situ*. On the other hand, it is theoretically possible that elevated levels of cytokines detected in vaginal fluid may also be the result of another phenomenon, such as vaginal transudation, a natural process that allows vaginal lubrication through vaso-dilatation.³⁴

A positive correlation between inflammatory cytokine levels and proviral load in the blood was described in HTLV-1-infected individuals diagnosed with Sicca syndrome.¹⁸ However, our study found no correlations between the levels of cytokines in vaginal fluid and proviral load in either cervicovaginal fluid or PBMCs. Of note, the HTLV-1-infected women evaluated herein were asymptomatic for diseases associated with this virus and had very low PVL in the vaginal fluid, 62 copies/10⁶ cells, about 0.006% of infected cells, in addition to intermediate levels of proviral load in the blood (28,665 copies/10⁶ cells, about 2.9% of infected PBMCs). For comparison, HTLV-1 PVL in peripheral blood is considered low when <1% of PBMCs are infected and intermediate only when 1–5% of PBMCs are infected.^{20,35}

With respect to the cytopathological findings and vaginal microbiota, no differences were observed between HTLV-1-infected and uninfected women. Higher levels of Th1 and Th2 cytokines were found in women with cervical intraepithelial neoplasia, with increasing levels seen in accordance with lesion severity.^{29,36} In our study, one woman from each group had atypical squamous cells of undetermined significance, a cytopathological alteration that requires periodic monitoring.

A limitation of the present study was the absence of HTLV-1-infected women diagnosed with HAM/TSP, who consistently present higher proviral loads in the peripheral blood as compared to asymptomatic individuals. However, HTLV-1 proviral load in the cervicovaginal fluid was indeed detectable in the majority of HTLV-1-asymptomatic women herein. Another limitation was that as cytokine levels in the plasma were not quantified, it was impossible to pair these with those in the cervical fluid.

Conclusions

The results presented herein show that HTLV-1 infection induces immune activation in the cervicovaginal environment of asymptomatic women, characterized by elevated levels of Th1, Th2, and IL17 cytokines in the cervicovaginal fluid. Further studies should be conducted involving HTLV-1-infected women who present high levels of proviral load in the peripheral blood to determine whether correlations exist regarding proviral load in cervicovaginal fluid.

Financial support

This work was supported by the Fundação de Amparo à Pesquisa do Estado da Bahia (BOL0575/2015). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) - Finance Code 001

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

We thank Mr. Noilson Lazaro and Viviana Olavarria for technical assistance. The authors would like to thank Andris K. Walter for English revision/editing services.

REFERENCES

- Gessain A, Gessain A, Cassar O. Epidemiological aspects and world distribution of HTLV-1 infection. *Front Microbiol.* 2012;3:388.
- Nunes D, Boa-Sorte N, Grassi MF, et al. HTLV-1 is predominantly sexually transmitted in Salvador, the city with the highest HTLV-1 prevalence in Brazil. *PLOS ONE.* 2017;12:e0171303. Epub 2017/02/06.
- Dourado I, Alcantara LC, Barreto ML, da Gloria Teixeira M, Galvao-Castro B. HTLV-I in the general population of Salvador, Brazil: a city with African ethnic and sociodemographic characteristics. *J Acquir Immune Defic Syndr.* 2003;34:527-31.
- Yoshida M, Miyoshi I, Hinuma Y. Isolation and characterization of retrovirus from cell lines of human adult T-cell leukemia and its implication in the disease. *Proc Natl Acad Sci U S A.* 1982;79:2031-5.
- Gessain A, Vernant JC, Maurs L, et al. Antibodies to human T-lymphotropic virus type-I in patients with tropical spastic paraparesis. *Lancet.* 1985;326:407-10.
- Mochizuki M, Watanabe T, Yamaguchi K, et al. Uveitis associated with human T lymphotropic virus type I: seroepidemiologic, clinical, and virologic studies. *J Infect Dis.* 1992;166:943-4. Epub 1992/10/01.
- La Grenade L. HTLV-I-associated infective dermatitis: past, present, and future. *J Acquir Immune Defic Syndr Hum Retrovirol.* 1996;13 Suppl. 1:S46-9. Epub 1996/01/01.
- Ferraz-Chaoui AK, Atta AM, Atta ML, Galvao-Castro B, Santiago MB. Study of autoantibodies in patients with keratoconjunctivitis sicca infected by the human T cell lymphotropic virus type 1. *Rheumatol Int.* 2010;30:775-8.
- Honarbaksh S, Taylor GP. High prevalence of bronchiectasis is linked to HTLV-1-associated inflammatory disease. *BMC Infect Dis.* 2015;15:258. Epub 2015/07/06.
- Nishioka K, Sumida T, Hasunuma T. Human T lymphotropic virus type I in arthropathy and autoimmune disorders. *Arthrit Rheumat.* 1996;39:1410-8.
- Popovic M, Lange-Wantzin G, Sarin PS, Mann D, Gallo RC. Transformation of human umbilical cord blood T cells by human T-cell leukemia/lymphoma virus. *Proc Natl Acad Sci U S A.* 1983;80:5402-6. Epub 1983/09/01.
- Richardson JH, Edwards AJ, Cruickshank JK, Rudge P, Dalgleish AG. In vivo cellular tropism of human T-cell leukemia virus type 1. *J Virol.* 1990;64:5682-7.
- Macatonia SE, Cruickshank JK, Rudge P, Knight SC. Dendritic cells from patients with tropical spastic paraparesis are infected with HTLV-1 and stimulate autologous lymphocyte proliferation. *AIDS Res Hum Retroviruses.* 1992;8:1699-706. Epub 1992/09/01.
- Carvalho EM, Bacellar O, Porto AF, Braga S, Galvao-Castro B, Neva F. Cytokine profile and immunomodulation in asymptomatic human T-lymphotropic virus type 1-infected blood donors. *J Acquir Immune Defic Syndr.* 2001;27:1-6.
- Chaves DG, Sales CC, de Cassia Goncalves P, et al. Plasmatic proinflammatory chemokines levels are tricky markers to monitoring HTLV-1 carriers. *J Med Virol.* 2016;88:1438-47. Epub 2016/01/24.
- Starling AL, Martins-Filho OA, Lambertucci JR, et al. Proviral load and the balance of serum cytokines in HTLV-1-asymptomatic infection and in HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP). *Acta Trop.* 2013;125:75-81.
- Nascimento MC, Primo J, Bittencourt A, et al. Infective dermatitis has similar immunological features to human T lymphotropic virus-type 1-associated myelopathy/tropical spastic paraparesis. *Clin Exp Immunol.* 2009;156:455-62. Epub 2009/05/15.
- Lima CM, Santos S, Dourado A, et al. Association of sicca syndrome with proviral load and proinflammatory cytokines in HTLV-1 infection. *J Immunol Res.* 2016;2016:8402059. Epub 2016/02/24.
- Castro-Lima Vargens C, Grassi MF, Boa-Sorte N, et al. Keratoconjunctivitis sicca of human T cell lymphotropic virus type 1 (HTLV-1) infected individuals is associated with high levels of HTLV-1 proviral load. *J Clin Virol.* 2011;52:177-80.
- Grassi MF, Olavarria VN, Kruschewsky Rde A, et al. cell lymphotropic virus type 1 (HTLV-1) proviral load of HTLV-associated myelopathy/tropical spastic paraparesis (HAM/TSP) patients according to new diagnostic criteria of HAM/TSP. *J Med Virol.* 2011;83:1269-74.
- Coutinho R Jr, Grassi MF, Korngold AB, et al. lymphotropic virus type 1 (HTLV-1) proviral load induces activation of T-lymphocytes in asymptomatic carriers. *BMC Infect Dis.* 2014;14:453. Epub 2014/08/26.
- Zunt JR, Dezzutti CS, Montano SM, et al. Cervical shedding of human T cell lymphotropic virus type I is associated with cervicitis. *J Infect Dis.* 2002;186:1669-72. Epub 2002/11/26.
- Belec L, Georges-Courbot MC, Georges A, et al. Cervicovaginal synthesis of IgG antibodies to the immunodominant 175-199 domain of the surface glycoprotein gp46 of human T-cell leukemia virus type I. *J Med Virol.* 1996;50:42-9. Epub 1996/09/01.
- Solomon D, Davey D, Kurman R, et al. The 2001 Bethesda System: terminology for reporting results of cervical cytology. *JAMA.* 2002;287:2114-9. Epub 2002/04/23.
- Dehee A, Cesaire R, Desire N, et al. Quantitation of HTLV-I proviral load by a TaqMan real-time PCR assay. *J Virol Methods.* 2002;102:37-51.
- Sato T, Coler-Reilly A, Utsunomiya A, Araya N, Yagishita N, Ando H. CSF CXCL10, CXCL9, and neopterin as candidate prognostic biomarkers for HTLV-1-associated myelopathy/tropical spastic paraparesis. *PLoS Negl Trop Dis.* 2013;7:e2479.
- Santos SB, Oliveira P, Luna T, et al. Immunological and viral features in patients with overactive bladder associated with human T-cell lymphotropic virus type 1 infection. *J Med Virol.* 2012;84:1809-17. Epub 2012/09/22.
- Fiorentino DF, Bond MW, Mosmann TR. Two types of mouse T helper cell. IV. Th2 clones secrete a factor that inhibits cytokine production by Th1 clones. *J Exp Med.* 1989;170:2081-95. Epub 1989/12/01.
- Tavares-Murta BM, de Resende AD, Cunha FQ, Murta EF. Local profile of cytokines and nitric oxide in patients with bacterial vaginosis and cervical intraepithelial neoplasia. *Eur J Obstet Gynecol Reprod Biol.* 2008;138:93-9. Epub 2007/08/09.
- Crowley-Nowick PA, Ellenberg JH, Vermund SH, Douglas SD, Holland CA, Moscicki AB. Cytokine profile in genital tract secretions from female adolescents: impact of human

- immunodeficiency virus, human papillomavirus, and other sexually transmitted pathogens. *J Infect Dis.* 2000;181:939-45. Epub 2000/03/18.
31. Belec L, Gherardi R, Payan C, et al. Proinflammatory cytokine expression in cervicovaginal secretions of normal and HIV-infected women. *Cytokine.* 1995;7:568-74. Epub 1995/08/01.
 32. Masson L, Salkinder AL, Olivier AJ, et al. Relationship between female genital tract infections, mucosal interleukin-17 production and local T helper type 17 cells. *Immunology.* 2015;146:557-67. Epub 2015/08/25.
 33. Zara F, Nappi RE, Brerra R, Migliavacca R, Maserati R, Spinillo A. Markers of local immunity in cervico-vaginal secretions of HIV infected women: implications for HIV shedding. *Sex Transm infect.* 2004;80:108-12. Epub 2004/04/01.
 34. Linhares IM, Giraldo PC, Baracat EC. Novos conhecimentos sobre a flora bacteriana vaginal. *J Rev Assoc Méd Bras.* 2010;56:370-4.
 35. Goncalves DU, Proietti FA, Barbosa-Stancioli EF, et al. HTLV-1-associated myelopathy/tropical spastic paraparesis (HAM/TSP) inflammatory network. *Inflamm Allergy Drug Targets.* 2008;7:98-107.
 36. Daniilidis A, Koutsos J, Oikonomou Z, Nasioutziki M, Hatziparadisi K, Tantanasis T. Cytokines of cervical mucosa and human papilloma virus infection of the cervix: a descriptive study. *Acta Cytol.* 2016;60:58-64. Epub 2016/ 03/24.