



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Brief communication

Trajectory of serogroups causing Invasive Meningococcal Disease in Santa Catarina state, Brazil (2007–2019)



Ricardo Becker Feijo ^{a,*}, Juarez Cunha ^b

^a Universidade Federal do Rio Grande do Sul, Faculdade de Medicina, Departamento de Pediatria, Porto Alegre, RS, Brazil

^b Secretaria Municipal de Saúde, Diretoria Geral de Vigilância em Saúde, Unidade de Vigilância Epidemiológica, Porto Alegre, RS, Brazil

ARTICLE INFO

Article history:

Received 29 September 2019

Accepted 13 June 2020

Available online 10 July 2020

Keywords:

Invasive meningococcal disease

Serogroup W

Santa Catarina state

Brazil

ABSTRACT

The aim of this study was to compare the trajectory of serogroups causing Invasive Meningococcal Disease (IMD) in the Santa Catarina (SC) state with those of whole Brazil. A retrospective analysis of all IMD cases reported from January 2007 to December 2019 was carried out. During the study period, 26,058 IMD cases were registered in Brazil and 644 in SC state alone. Overall, Brazil showed progressive reduction in cases since 2010, when the meningococcal C conjugate vaccine was introduced on National Immunization Program, while SC showed an increase in total cases since 2013, particularly from serogroups W and C. Serogroups distribution was significantly different between Brazil and SC. The emergence of serogroup W highlights the improved meningococcal surveillance through increased accuracy in identification methods in SC state. This finding is important for discussing recommendations of quadrivalent (ACWY) conjugate vaccines in different geographical areas of Brazil.

© 2020 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Neisseria meningitidis (*N. meningitidis*) causes extremely severe conditions in humans, especially invasive meningococcal disease (IMD) characterized by one or more clinical syndromes including bacteremia, sepsis, or meningitis, the latter being the most common presentation.¹

Capsular groups A, B, C, W, Y, and X cause almost all IMD.² However, the quality and reliability of the information on IMD are not uniform across the world, due in part to differences in surveillance practices, use of different diagnostic methods and protocols, and application of different meningococcal case definition.³ Since late 1960s capsular group W has become a

frequent cause of IMD,⁴ particularly in Europe, South America, Australia, and some parts of Sub-Saharan Africa.⁵

The highest incidence rates of IMD are observed in children aged less than one year,^{6,7} but in some regions other incidence peaks are observed among adolescents (16 through 21 years of age)^{8,9} and those aged ≥65 years.^{10,11} In Latin America, the incidence of IMD varies from <0.1 cases per 100,000 in countries including Mexico, Paraguay, Peru, and Bolivia to almost two cases per 100,000 in Brazil.⁷ In Brazil, *N. meningitidis* was the primary etiology of acute bacterial meningitis with serogroup C being the most prevalent in the country. In

* Corresponding author at: Rua Attilio Bilibio 120/23, 91530008 Porto Alegre, RS, Brazil.

E-mail address: rfeijo@hcpa.edu.br (R.B. Feijo).

<https://doi.org/10.1016/j.bjid.2020.06.007>

1413-8670/© 2020 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 – Meningococcal Disease in Brazil compared to Santa Catarina State. NT, nontypeable; OTHER, Other serogroups.

Table 1 – Meningococcal serogroup distribution during the period of 2007-2019.

	Serogroups					Total
	NT ^a	B	C	W	OTHER ^b	
Brazil	13,589 (52.1%)	2608 (10.0%)	8728 (33.5%)	802 (3.1%)	331 (1.3%)	26,058 (100%)
SC ^c	291 (45.2%)	106 (16.5%)	136 (21.1%)	101 (15.7%)	10 (1.6%)	644 (100%)

^a Nontypeable.

^b Other serogroups.

^c Santa Catarina State.

2010, the National Immunization Program (NIP) included the MenC conjugate vaccine for children under two years old, with no catch-up campaign in older ages groups. Based on the significant IMD reduction observed only in the vaccinated group, the Ministry of Health decided to include MenC vaccine for adolescents aged 11–14 years in 2017.¹²

The introduction of multiplex real-time PCR (polymerase chain reaction) (RT-PCR) assay testing for *S. pneumoniae*, *N. meningitidis* and *H. influenzae* type B in sentinel hospitals in São Paulo increased the diagnostic yield for bacterial meningitis by 52, 85 and 20%, respectively, over culture-based methods.¹³ Since then, there has been a progressive increase in the use of PCR assays with improvement in microbiologic confirmation of bacterial meningitis throughout the country.

IMD is associated with substantial morbidity and overall case fatality rates (CFRs) of around 10% (with CFRs as high as 20% reported in some countries).¹⁴ Meningococcal disease caused by serogroup W alone has been associated with CFRs of >30%.¹⁵

Serogroups B and C are currently dominant in Latin America. Nevertheless, the proportion of IMD cases including outbreaks attributable to other serogroups have been reported in several countries.¹⁶ Southern Brazil, Argentina and Chile witnessed the emergence of MenW cc11 strain as a major cause of endemic meningococcal disease after 2003.¹⁷ The states of Paraná, Santa Catarina and Rio Grande do Sul in the southern region of Brazil borders the southern cone countries of Latin America. In fact, W serogroup represented 17.8% of all invasive strains in 2003–2005 compared to 3.2% in 1995–2002.¹⁸

Notifications of IMD in Brazil are routinely collected by the Minister of Health through the Information System for Notifiable Diseases (SINAN). This study is a retrospective review of all IMD cases reported in Brazil as a whole and in SC in particular, from 1 January 2007 to 31 December 2019, using the SINAN electronic database.^{19,20}

The study variables included the number total cases of IMD and meningococcal serogroups. Pearson's chi squared test was used to compare total cases and serogroups frequencies. For comparing distributions evolution, stacked bars were used as visualization. Statistical software used was R version 3.5.2.

Total number of cases and cases according to serogroups were different between Brazil and Santa Catarina state along the reported years (Fig. 1). Analysis of total number of cases by year from 2007 through 2019 showed statistically significant differences between Brazil and SC state ($p < 0.001$) (Table 1).

A progressive decline of total cases and its respective serogroups occurred after introduction of meningococcal C vaccination in 2010 in Brazil. In contrast, there was a diminishing number of cases in SC state from 2007 to 2013, with a reversal of trends from 2014 to 2018, with a reduction of the number of the cases in 2019.

Serogroup distribution showed specific changes in SC state. Meningococcal disease by serogroup W increased between 2016–2018, with a reduction in 2019. Unlike Brazil, serogroup C showed an increment of cases after 2013 in SC state.

In general, non-identified cases showed decreasing trends in both regions even though they have occurred in different proportions.

Differences between the number of cases and distribution of serogroups cannot be clearly explained. There are several variables that can explain the changes in the prevalence of individual serogroups, one of them could be the naturally occurring periodic changes and another the impact of immunization programs. Another possibility, considering that SC is a region with high affluence of tourism (mainly from Argentina), is that these differences could be influenced by tourism carriage.

The notification of SC state presented a cut point at December 2nd, 2019, which might affect the analysis of the following months of the analyzed period.

The observed number of cases and serogroup distribution along time for SC state compared to Brazil, could imply that health authorities need to review current policies in order to cope with existing epidemiology, especially since the W serogroup is associated directly with increased mortality and morbidity rates.^{15,16}

The emergence of serogroup W in SC highlights the difference in how health authorities have been dealing with meningococcal surveillance as the identification methods increased in accuracy. Moreover, it shows that geographical features should be considered as an important variable when recommending quadrivalent (ACWY) conjugate vaccines exceptionally considering the emergency of a strain that used to be underrepresented.

In summary, we showed that the total number of cases and serogroup distribution in the time period of 2007 through 2019 in SC state was significantly different compared to Brazil.

Funding

Sponsorship of this study and article processing charges were funded by Pfizer Inc.

Conflict of interest

The authors are consultant and speaker Pfizer, Sanofi, Glaxo-SmithKline and Merck Sharp & Dohme.

Acknowledgments

Sponsorship of this study and article processing charges were funded by Pfizer Inc. All named authors meet the International Committee of Medical Journal Editors criteria for authorship for this article, take responsibility for the integrity of the work as a whole, and have given their approval for this version to be published. The statistical analysis was conducted by Content Medicine in Buenos Aires, Argentina; funding was provided by Pfizer Inc. This article is based on previously conducted studies and does not contain any studies with human participants or animals performed by any of the authors.

REFERENCES

1. Dwilow R, Fanella S. Invasive meningococcal disease in the 21st century—an update for the clinician. *Curr Neurol Neurosci Rep.* 2015;15:2.
2. Harrison OB, Claus H, Jiang Y, Bennett JS, Bratcher HB, Jolley KA, et al. Description and nomenclature of *Neisseria meningitidis* capsule locus. *Emerg Infect Dis.* 2013;19:566–73.
3. Sáfadi MAP, McIntosh EDG. Epidemiology and prevention of meningococcal disease: a critical appraisal of vaccine policies. *Expert Rev Vaccines.* 2011;10:1717–30.
4. Araya P, Fernández J, Del Canto F, Seoane M, Ibarz-Pavón AB, Barra G, et al. *Neisseria meningitidis* ST-11 clonal complex, Chile 2012. *Emerg Infect Dis.* 2015;21:339–41.
5. Harrison LH. Epidemiological profile of meningococcal disease in the United States. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 2010;50:S37–44.
6. Harrison LH, Pelton SI, Wilder-Smith A, Holst J, Safadi MAP, Vazquez JA, et al. The Global Meningococcal Initiative: recommendations for reducing the global burden of meningococcal disease. *Vaccine.* 2011;29:3363–71.
7. Sáfadi MAP, Cintra OAL. Epidemiology of meningococcal disease in Latin America: current situation and opportunities for prevention. *Neurol Res.* 2010;32:263–71.
8. National Advisory Committee on Immunization (NACI). An update on the invasive meningococcal disease and meningococcal vaccine conjugate recommendations. An Advisory Committee Statement (ACS). *Can Commun Dis Rep Relevé Mal Transm Au Can.* 2009;35:1–40.
9. Cohn AC, MacNeil JR, Harrison LH, Hatcher C, Theodore J, Schmidt M, et al. Changes in *Neisseria meningitidis* disease epidemiology in the United States, 1998–2007: implications for prevention of meningococcal disease. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 2010;50:184–91.
10. Abad R, Agudelo CI, Brandileone MC, Chanto G, Gabastou JM, Hormazabal JC, et al. Molecular characterization of invasive serogroup Y *Neisseria meningitidis* strains isolated in the Latin America region. *J Infect.* 2009;59:104–14.
11. Cohn AC, MacNeil JR, Clark TA, Ortega-Sánchez IR, Briere EZ, Meissner HC, et al. Prevention and control of meningococcal disease: recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR Recomm Rep Morb Mortal Wkly Rep Recomm Rep.* 2013;62:1–28.
12. Sáfadi MAP, Gonzalez-Ayala S, Jakel A, Wieffer H, Moreno C, Vyse A. The epidemiology of meningococcal disease in Latin America 1945–2010: an unpredictable and changing landscape. *Epidemiol Infect.* 2013;141:447–58.
13. Sacchi CT, Fukasawa LO, Gonçalves MG, Salgado MM, Shutt KA, Carvalhanas TR, et al. Incorporation of Real-Time PCR into Routine Public Health Surveillance of Culture Negative Bacterial Meningitis in São Paulo, Brazil. *PLOS ONE.* 2011;6:e20675.
14. Erickson L, De Wals P. Complications and sequelae of meningococcal disease in Quebec, Canada, 1990–1994. *Clin Infect Dis Off Publ Infect Dis Soc Am.* 1998;26:1159–64.
15. Moreno G, López D, Vergara N, Gallegos D, Advis MF, Loayza S. Clinical characterization of cases with meningococcal disease by W135 group in Chile, 2012. *Rev Chil Infectol.* 2013;30:346–9.
16. Valenzuela MT, Moreno G, Vaquero A, Seoane M, Hormazábal JC, Bertoglia MP, et al. [Emergence of W135 meningococcal serogroup in Chile during 2012]. *Rev Med Chil.* 2013;141:959–67.
17. Abad R, López EL, Debbag R, Vázquez JA. Serogroup W meningococcal disease: global spread and current affect on the Southern Cone in Latin America. *Epidemiol Infect.* 2014;142:2461–70.
18. Weidlich L, Baethgen LF, Mayer LW, Moraes C, Klein CC, Nunes LS, et al. High prevalence of *Neisseria meningitidis* hypervirulent lineages and emergence of W135:P1.5,2:ST-11 clone in Southern Brazil. *J Infect.* 2008;57:324–31.
19. DATASUS [Internet]. [cited 2020 Mar 9]. Available from: <http://www2.datasus.gov.br/DATASUS/index.php?area=0203&id=29892234&VObj>.
20. DIVE - Boletim Epidemiológico Mensal – Vigilância da Doença Meningocócica [Internet]. [cited 2020 Mar 9]. Available from: <http://www.dive.sc.gov.br/index.php/arquivo-noticias/805-boletim-epidemiologico-mensal-n-10-2018-vigilancia-da-doenca-meningococica-atualizado-em-04-de-janeiro-de-2019>.