



The Brazilian Journal of INFECTIOUS DISEASES

www.elsevier.com/locate/bjid



Original Article

SARS-CoV-2 testing disparities across geographical regions from a large metropolitan area in Brazil: Results from a web-based survey among individuals interested in clinical trials for COVID-19 vaccines

Thiago S. Torres ^{a,*}, Paula M. Luz ^a, Lara E. Coelho ^a, Cristina Jalil ^a, Gisely G. Falco ^a, Leonardo P. Sousa ^a, Emilia Jalil ^a, Daniel R.B. Bezerra ^a, Sandra W. Cardoso ^a, Brenda Hoagland ^a, Claudio J. Struchiner ^b, Valdilea G. Veloso ^a, Beatriz Grinsztejn ^a

^a Instituto Nacional de Infectologia Evandro Chagas, Fundação Oswaldo Cruz, Rio de Janeiro, RJ, Brazil

^b Fundação Getúlio Vargas (FGV), Rio de Janeiro, RJ, Brazil

ARTICLE INFO

Article history:

Received 18 May 2021

Accepted 18 July 2021

Available online 2 August 2021

Keywords:

SARS-CoV-2

COVID-19

Brazil

Seroprevalence

Vaccine

Survey

ABSTRACT

After more than a year since the novel coronavirus (SARS-CoV-2) disease 2019 or COVID-19 has reached the status of a global pandemic, the number of COVID-19 cases continues to rise in Brazil. As no effective treatment been approved yet, only mass vaccination can stop the spread of SARS-CoV-2 and end the COVID-19 pandemic. Multiple COVID-19 vaccine candidates are under development and some are currently in use. This study aims to describe the characteristics of individuals who have registered in an online platform to participate in clinical trials for COVID-19 vaccines. Additionally, participants' characteristics according to age and presence of comorbidities associated with severe COVID-19 and differences of SARS-CoV-2 testing across different geographical areas/neighborhoods are provided. This was a cross-sectional web-based study conducted between September and December/2020, aiming to reach individuals aged ≥ 18 years who live in Rio de Janeiro metropolitan area, Brazil. Among 21,210 individuals who completed the survey, 20,587 (97.1%) were willing to participate in clinical trials for COVID-19 vaccines. Among those willing to participate, 57.8% individuals were aged 18–59 years and had no comorbidity, 33.7% were aged 18–59 years and had at least one comorbidity, and 8.6% were aged ≥ 60 years regardless the presence of any comorbidity. Almost half (42.6%) reported ever testing for COVID-19, and this proportion was lower among those aged ≥ 60 years ($p < 0.001$). Prevalence of positive PCR results was 16.0%, higher among those aged 18–59 years ($p < 0.009$). Prevalence of positive antibody result was 10.0%, with no difference across age and comorbidity groups. Participants from areas/neighborhoods with higher Human Development Index (HDI) reported ever testing for SARS-CoV-2 more frequently than those from lower HDI areas. Interest to participate in clinical trials for COVID-19 vaccines candidates in Rio de

* Corresponding author.

E-mail address: thiago.torres@ini.fiocruz.br (T.S. Torres).

<https://doi.org/10.1016/j.bjid.2021.101600>

1413-8670/© 2021 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Janeiro was significantly high. The online registry successfully reached out a large number of individuals with diverse sociodemographic, economic and clinical backgrounds.

© 2021 Sociedade Brasileira de Infectologia. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Introduction

The novel coronavirus (SARS-CoV-2) disease 2019, or COVID-19, is currently the largest and most challenging public health crisis worldwide, having reached the status of a global pandemic since March 11, 2020.¹ After more than a year, the number of cases of COVID-19 continues to rise in Brazil. From February 26, 2020 to August 10, 2021 there have been 20,212,642 confirmed cases and more than 564,000 deaths in the country. The Southeast region, which includes the Rio de Janeiro State, accounted for 38% of all COVID-19 cases in Brazil. Rio de Janeiro State confirmed 1,063,772 cases, holding the lowest incidence rate per 100,000 inhabitants, but the highest mortality rate in the region (348.3 deaths/100,000 inhabitants).²

Since the onset of the pandemic, social distancing and community containment measures to avoid the spread of COVID-19 and the collapse of the health system have been poorly and unequally adopted in the country.³ During May 2021, only 30% of Brazilians reported being isolated at home, the lowest rate since the onset of the pandemic.⁴ Despite the major importance of such measures, low income Brazilians struggle to adhere to social distancing measures due to financial constraints.^{5,6} Brazil is the 9th largest economy by nominal Gross Domestic Product (GDP = US\$ 1.9 trillion), but the 9th most unequal country in the world.⁷ Brazilian geographic regions have impressive disparities in terms of GDP and Human Development Index (HDI), and these disparities are dramatic when analyzed within cities and metropolitan areas. Rio de Janeiro metropolitan area is the second largest in the country and the 16th of the world, accounting for more than 13 million inhabitants and 22 municipalities. Huge socioeconomic disparities are widespread in the region, in which HDI varies from very high to low in adjacent neighborhoods.⁸ These inequalities in wealth and health care closely impact health outcomes,⁹ including hospitalizations and deaths related to COVID-19.¹⁰

Mass vaccination campaigns to prevent or respond to outbreaks of vaccine-preventable diseases are effective strategies to reduce the number of cases and deaths.¹¹ With no effective treatment currently approved, only mass vaccination may stop the spread of SARS-CoV-2 and end COVID-19 pandemic. By May 2021, four COVID-19 vaccines have been registered or approved for emergency use by the Brazilian Health Regulatory Agency (ANVISA). The ChAdOx1 nCoV-19,¹² developed by AstraZeneca/Oxford and produced by Bio-Manguinhos, Fundação Oswaldo Cruz (Fiocruz),¹³ and the BNT162b2,¹⁴ produced by Pfizer, Inc. and BioNTech have been registered.¹⁵ CoronaVac,¹⁶ developed by Sinovac Life Science Co. Ltd. and produced by Instituto Butantan,¹⁷ and Ad26.COV2.S, developed and produced by Janssen Vaccines & Prevention B.V.¹⁸

have been approved for the emergency use only. From January 17, 2021 to August 10, 2021, 109,208,435 Brazilians have received at least one dose of COVID-19 vaccine and 46,902,420 were fully vaccinated, representing 22.1% of the country's population.¹⁹

Additional COVID-19 vaccine candidates are under development and some will get to clinical development soon.²⁰ Due to the high prevalence of COVID-19 in Brazil, clinical trials may be developed in the country and research centers should be structured and prepared to conduct these trials. Instituto Nacional de Infectologia Evandro Chagas (INI)-Fiocruz, located in Rio de Janeiro, Brazil, is a centennial institution fully dedicated to research, prevention and care of infectious diseases, including HIV, tuberculosis, Zika, Dengue, and hepatitis, with a long path of conducting large clinical trials.²¹ With the advent of the COVID 19 pandemic, the Institute has fully embraced the mission of providing high level care and research on this new deadly disease.

This study aims to describe the characteristics of individuals living in Rio de Janeiro metropolitan area who have registered in an online platform to participate in clinical trials for COVID-19 vaccines candidates to be conducted at INI-Fiocruz. Additionally, we compare participants' characteristics according to age and presence of comorbidities associated with severe COVID-19 and provide differences of SARS-CoV-2 testing across different areas/neighborhoods of the area.

Material and methods

Study design

This was a cross-sectional web-based study, in which participants were invited to complete an online questionnaire and provide personal information such as name, mobile phone number and e-mail, to be potentially contacted by INI-Fiocruz team for COVID-19 vaccines clinical trials. Eligibility criteria were age ≥ 18 years and residence in Rio de Janeiro metropolitan area. Individuals were excluded from this analysis if they had previously completed the questionnaire (same mobile phone number), did not inform neighborhood of domicile, reported being pregnant, or reported no interest to participate in clinical trials for COVID-19 vaccines.

The online questionnaire was programmed on Alchemer®. The researchers piloted the questionnaire before starting the study. The survey was open (not password-protected). Mobile version of the survey had one question per page, and number of pages depended on the chosen answers (adaptive questioning). All survey items had a non-response option ("I don't want to answer"). CHERRIES statement for web-based survey was followed.²²

The survey was launched on September 10, 2020. On September 23, 2020 it was advertised on INI-Fiocruz (<https://www.ini.fiocruz.br/>) and Fiocruz webpages (<https://portal.fiocruz.br/noticia/fiocruz-cadastra-voluntarios-para-participar-de-estudo-de-vacina-para-prevencao-da-covid-19>). We also implemented community advertisement in low income areas of Rio de Janeiro city (INI-Fiocruz neighbor communities and West region) using posters, banners and flyers with telephone number, WhatsApp contact and QR code, which directed individuals to the survey link. We identified micro influencers at these regions, who performed lectures and meetings to explain the importance of COVID-19 vaccines. To increase the percentage of answers among individuals aged 55+ years, we posted advertisements on Facebook from November 18 to 23, 2020 targeting this population. We closed the survey link and stopped recruitment on December 07, 2020.

The INI-Fiocruz institutional review board (#CAAE 36868820.9.0000.5262) reviewed and approved this study. All participants acknowledged reading the informed consent text and provided digital consent. No incentives were provided in accordance with Brazilian regulations.

Survey instrument

The survey instrument was composed of six sections (33 questions) addressing: sociodemographic, daily activities, comorbidities, and SARS-CoV-2 testing.

Variables

Sociodemographic

Variables were age at the time of the survey; gender identity (cisgender men, cisgender women or transgender/non-binary); region (city and neighborhood); race/color (Asian, Black, Indigenous, *Pardo*, White, and "I don't know"); schooling (according to Brazilian educational levels, from elementary incomplete to graduation); and number of people living at home (alone, 1, 2, 3, 4, or ≥ 5).

Daily activities

Participants were inquired about current daily activities such as: receiving visits at home, which included people in charge of services, such as cleaning (never, rarely, monthly, weekly, or daily), and working (yes, retired, unemployed, or license). Those reporting current work were asked about frequency of commuting to work (never, once a week, 2 to 4 times a week or 5, or more times a week). Those commuting to work at least once a week answered questions about means of transportation to work and use of personnel protective equipment (PPE) such as face mask by coworkers (yes, no, or I don't know). In addition, we inquired about participation in public events with 10+ people in the last two weeks (yes, no, or I don't know).

Associated factors with severe COVID-19

Participants were asked about current or past comorbidities or conditions that are associated with severe COVID-19 disease from a pre-determined list: any cardiovascular disease (e.g. heart failure, coronary artery disease, congenital heart disease, cardiomyopathies, and pulmonary hypertension),

sleep apnea, arthritis, asthma or bronchitis, stroke (ischemic, hemorrhagic, or intracranial), cancer, neurological conditions, diabetes (type I or II), chronic liver disease (e.g. steatosis, cirrhosis, chronic hepatitis B), chronic kidney disease (requires hemodialysis or peritoneal dialysis), COPD (chronic obstructive pulmonary disease or pulmonary emphysema), pulmonary fibrosis or cystic fibrosis, systemic arterial hypertension, HIV, obesity (body mass index [BMI] ≥ 30), thalassemia or sickle cell anemia, active or treated tuberculosis (TB), and organ transplantation. Number of comorbidities was categorized into none, one, two, and three or more. Participants also provided information on tobacco/e-cigarette smoking (never, former, or current smoker).

SARS-CoV-2 testing and results

Participants were asked if ever tested for SARS-CoV-2 (yes/no). For those who answered positively, we asked which was the test (PCR [polymerase chain reaction] or antibody tests) and their results (positive, negative, inconclusive, or pending).

Statistical analysis

Sociodemographic characteristics, daily activities and SARS-CoV-2 testing were described for the entire sample and stratified into three mutually exclusive groups depending on the risk of severe COVID-19 disease (18–59 years with no comorbidity, 18–59 years with at least one comorbidity, ≥ 60 years regardless of comorbidities). Number and type of comorbidities were described according to age (< 60 years vs. ≥ 60 years). We compared SARS-CoV-2 testing and results as well as comorbidities among the three groups using chi-square test. Ever testing for SARS-CoV-2, PCR and antibody test results were described according to metropolitan area municipalities²³ and Rio de Janeiro city administrative region (33 regions)^{24,25} classified according to HDI.²⁶ Analyses were performed using Software R version 4.0.3.²⁷

Results

A total of 35,374 individuals accessed the questionnaire; 4197 (11.9%) on the day of advertisement was launched at Fiocruz website (23-Sep-2020), and 16,121 (45.6%) during the following seven days (24 to 30-Sep-2020). Among 21,210 individuals who completed the survey, only 623 (2.9%) reported no interest to participate in clinical trials for COVID-19 vaccines; the remaining 20,587 individuals (97.1%) were included in this analysis (Fig. A).

Median age was 36 years (interquartile range [IQR]: 26–47) and 1763 (8.6%) were aged ≥ 60 years. More than half were cisgender women (56.4%), 12,457 (60.5%) were white, 11,452 (55.6%) reported completed college education or higher, 5123 (24.9%) reported living with 5+ persons and only 3047 (14.8%) were not receiving visits at home (Table 1). Of all respondents, 30.1% attended a public event with 10 or more people in the previous two weeks. A total of 5334 (25.9%) were unemployed and 13,601 (66.1%) were currently working; 64.9% (8,832/13,601) reported commuting to work, and 41.1% (3,619/8,792) were using bus, train, metro or ferry as the main means of




Fig. A – Study flow-chart. Brazil, 2020.

transportation. The majority reported use of PPE at work (84.8%; 7488/8,832).

Overall, 8153 (39.6%) reported at least one comorbidity and 1038 (5.0%) reported three or more comorbidities (Table 2). Only 551 (31.3%) participants aged ≥ 60 years reported no comorbidity, and among this age group, 281 (15.9%) reported three or more comorbidities ($p < .001$). Overall, the most frequent comorbidities were obesity (14.7%), hypertension (14.0%), and asthma (9.0%). Among participants aged ≥ 60 years, the most frequent were hypertension (42.2%),

diabetes (18.6%), and obesity (13.6%). Only 2520 (12.2%) of the respondents reported being a current smoker and 4567 (22.2%) were former smokers.

Considering the three mutually exclusive groups, 11,883 (57.8%) were aged 18–59 years and reported no comorbidity, 6941 (33.7%) were aged 18–59 years and with at least one reported comorbidity, and 1763 (8.6%) were aged ≥ 60 years regardless of any comorbidity. Comparing these three groups, we noted that participants aged ≥ 60 years ($N = 1763$) were more likely to live in Rio de Janeiro city (including the most

Table 1 – Sociodemographic and daily activities characteristics of study population stratified by age and comorbidity. Rio de Janeiro, Brazil, 2020.

	Total	18–59 Years		≥ 60 years
		No comorbidity	At least one comorbidity	
	20,587	11,883 (57.8)	6941 (33.7)	1763 (8.6)
Gender				
Cisgender men	8885 (43.2)	4860 (40.9)	3196 (46.0)	829 (47.0)
Cisgender women	11,605 (56.4)	6985 (58.8)	3691 (53.2)	929 (52.7)
Transgender/non-binary	97 (0.5)	38 (0.3)	54 (0.8)	5 (0.3)
Region ¹				
Rio de Janeiro city	15,981 (77.6)	9221 (77.6)	5391 (77.7)	1441 (81.7)
Zona Norte	3261 (15.8)	1840 (15.5)	1194 (17.2)	227 (12.9)
Zona Sul	3018 (14.7)	1757 (14.8)	832 (12.0)	429 (24.3)
Barra da Tijuca and Jacarepaguá	2719 (13.2)	1597 (13.4)	852 (12.3)	270 (15.3)
Grande Tijuca	2105 (10.2)	1247 (10.5)	672 (9.7)	186 (10.6)
Zona Oeste	1747 (8.5)	945 (8)	692 (10)	110 (6.2)
Grande Meier	1418 (6.9)	801 (6.7)	505 (7.3)	112 (6.4)
Centro	969 (4.7)	565 (4.8)	340 (4.9)	64 (3.6)
Ilha do Governador	744 (3.6)	469 (3.9)	232 (3.3)	43 (2.4)
Baixada Fluminense	2490 (12.1)	1442 (12.1)	890 (12.8)	158 (9.0)
Niterói	1206 (5.9)	707 (5.9)	379 (5.5)	120 (6.8)
Leste Fluminense	766 (3.7)	437 (3.7)	296 (4.3)	33 (1.9)

Table 1 (continued)

	Total	18–59 Years		≥ 60 years
		No comorbidity	At least one comorbidity	
	20,587	11,883 (57.8)	6941 (33.7)	1763 (8.6)
Petropolis	144 (0.7)	76 (0.6)	57 (0.8)	11 (0.6)
Race				
White	12457 (60.5)	7152 (60.2)	4018 (57.9)	1287 (73.0)
Pardo	5334 (25.9)	3055 (25.7)	1925 (27.7)	354 (20.1)
Black	2330 (11.3)	1383 (11.6)	850 (12.2)	97 (5.5)
Indigenous	61 (0.3)	31 (0.3)	24 (0.3)	6 (0.3)
Asian	80 (0.4)	60 (0.5)	17 (0.2)	3 (0.2)
Don't know	325 (1.6)	202 (1.7)	107 (1.5)	16 (0.9)
Education				
Elementary incompleted	281 (1.4)	75 (0.6)	123 (1.8)	83 (4.7)
Elementary completed	364 (1.8)	149 (1.3)	135 (1.9)	80 (4.5)
Secondary incompleted	511 (2.5)	259 (2.2)	191 (2.8)	61 (3.5)
Secondary completed	2667 (13.0)	1418 (11.9)	977 (14.1)	272 (15.4)
Superior incompleted	5290 (25.7)	3478 (29.3)	1615 (23.3)	197 (11.2)
Superior completed	5380 (26.1)	2987 (25.1)	1824 (26.3)	569 (32.3)
Graduation	6072 (29.5)	3507 (29.5)	2067 (29.8)	498 (28.2)
Don't know	22 (0.1)	10 (0.1)	9 (0.1)	3 (0.2)
Number of people living at home				
Alone	3112 (15.1)	1632 (13.7)	1023 (14.7)	457 (25.9)
1	2504 (12.2)	1428 (12.0)	751 (10.8)	325 (18.4)
2	4239 (20.6)	2494 (21.0)	1363 (19.6)	382 (21.7)
3	2983 (14.5)	1747 (14.7)	1031 (14.9)	205 (11.6)
4	2626 (12.8)	1565 (13.2)	929 (13.4)	132 (7.5)
≥ 5	5123 (24.9)	3017 (25.4)	1844 (26.6)	262 (14.9)
Visit at home				
Never	3047 (14.8)	1782 (15.0)	1013 (14.6)	252 (14.3)
Rarely	8558 (41.6)	4892 (41.2)	2964 (42.7)	702 (39.8)
Monthly	1776 (8.6)	1108 (9.3)	588 (8.5)	80 (4.5)
Weekly	5747 (27.9)	3335 (28.1)	1878 (27.1)	534 (30.3)
Daily	1458 (7.1)	765 (6.4)	498 (7.2)	195 (11.1)
Current working				
Yes	13,601 (66.1)	8091 (68.1)	4787 (69.0)	723 (41.0)
Retired	1263 (6.1)	137 (1.2)	242 (3.5)	884 (50.1)
Unemployed	5334 (25.9)	3492 (29.4)	1730 (24.9)	112 (6.4)
Leave	389 (1.9)	163 (1.4)	182 (2.6)	44 (2.5)
Commute to work (N = 13,601)				
No	4769 (35.1)	2781 (34.4)	1685 (35.2)	303 (41.9)
Once a week	1171 (8.6)	676 (8.4)	417 (8.7)	78 (10.8)
2 to 4 times a week	3483 (25.6)	2128 (26.3)	1172 (24.5)	183 (25.3)
5+ times a week	4178 (30.7)	2506 (31.0)	1513 (31.6)	159 (22)
Main means of transportation to work (N = 8792)				
Walking or cycling	791 (9.0)	485 (9.1)	263 (8.5)	43 (10.2)
Ferry	66 (0.7)	40 (0.8)	22 (0.7)	4 (1)
Own car	3121 (35.3)	1768 (33.3)	1126 (36.3)	227 (54.0)
Car ride	291 (3.3)	187 (3.5)	95 (3.1)	9 (2.1)
Bus	2555 (29.1)	1611 (30.3)	892 (28.8)	52 (12.4)
Taxi or transport through apps	880 (10.0)	513 (9.7)	318 (10.3)	49 (11.7)
Train or metro	998 (11.3)	624 (11.7)	342 (11.0)	32 (7.6)
Mototaxi	90 (1.0)	59 (1.1)	29 (0.9)	2 (0.5)
PPE use at work (N = 8,832)				
Yes	7488 (84.8)	4517 (85.1)	2598 (83.8)	373 (88.8)
No	981 (11.1)	592 (11.1)	358 (11.5)	31 (7.4)
Don't know	361 (4.1)	201 (3.8)	144 (4.6)	16 (3.8)
Public events with 10+ people				
No	14,263 (69.3)	8245 (69.4)	4623 (66.6)	1395 (79.1)
Yes	6195 (30.1)	3555 (29.9)	2278 (32.8)	362 (20.5)
Don't know	128 (0.6)	82 (0.7)	40 (0.6)	6 (0.3)

1: Rio de Janeiro city 33 administrative regions were grouped into: Zona Norte (Ramos, Penha, Inhaúma, Irajá, Madureira, Anchieta, Pavuna, Jacarezinho, Complexo do Alemão, Maré, Vigário Geral), Zona Sul (Botafogo, Copacabana, Lagoa, Rocinha), Barra da Tijuca and Jacarepaguá (Jacarepaguá, Barra da Tijuca, Cidade de Deus), Grande Tijuca (Tijuca, Vila Isabel), Zona Oeste (Bangu, Campo Grande, Santa Cruz, Guaratiba, Realengo), Grande Méier, Centro (Zona Portuária, Centro, Rio Comprido, Santa teresa, São Cristóvão, Ilha de Paqueta), Ilha do Governador. Other municipalities of metropolitan area in: Baixada Fluminense (Duque de Caxias, Guapimirim, Itaguaí, Japeri, Magé, Mesquita, Nilópolis, Nova Iguaçu, Paracambi, Seropédica); Niterói; Leste Fluminense (Cachoeiras de Macacu, Itaboraí, Maricá, Rio Bonito, São Gonçalo, Tanguá); Petropolis. PPE: Personal Protection Equipment.

Table 2 – Factors associated with the risk of developing severe COVID-19 (self-reported) among study participants according to age. Rio de Janeiro, Brazil, 2020.

	Total N = 20,587	18–59 Years N = 18,824	≥ 60 Years N = 1763	p-value
Number of comorbidities				
Mean (SD)	0.6 (0.9)	0.5 (0.9)	1.3 (1.3)	< .001
None	12434 (60.4)	11883 (63.1)	551 (31.3)	< .001
1	5254 (25.5)	4661 (24.8)	593 (33.6)	
2	1861 (9.0)	1523 (8.1)	338 (19.2)	
3+	1038 (5.0)	757 (4.0)	281 (15.9)	
Type of Comorbidity				
Obesity	3023 (14.7)	2784 (14.8)	239 (13.6)	0.16
Hypertension	2873 (14.0)	2129 (11.3)	744 (42.2)	< 0.001
Asthma or bronchitis	1845 (9.0)	1734 (9.2)	111 (6.3)	< 0.001
Diabetes (type I or II)	1091 (5.3)	763 (4.1)	328 (18.6)	< 0.001
Apnea	958 (4.7)	823 (4.4)	135 (7.7)	< 0.001
Cardiovascular diseases	566 (2.7)	385 (2.0)	181 (10.3)	< 0.001
HIV	512 (2.5)	487 (2.6)	25 (1.4)	0.003
Arthritis	461 (2.2)	300 (1.6)	161 (9.1)	< 0.001
Chronic liver disease	376 (1.8)	298 (1.6)	78 (4.4)	< 0.001
Cancer	251 (1.2)	136 (0.7)	115 (6.5)	< 0.001
Treated TB	156 (0.8)	126 (0.7)	30 (1.7)	< 0.001
COPD	137 (0.7)	74 (0.4)	63 (3.6)	< 0.001
Neurological conditions	127 (0.6)	110 (0.6)	17 (1.0)	0.051
Thalassemia	94 (0.5)	90 (0.5)	4 (0.2)	0.13
Stroke	58 (0.3)	34 (0.2)	24 (1.4)	< 0.001
Pulmonary fibrosis	22 (0.1)	12 (0.1)	10 (0.6)	< 0.001
Chronic kidney disease	19 (0.1)	15 (0.1)	4 (0.2)	0.052
Active TB	9 (0)	9 (0)	0 (0)	0.34
Organ transplantation	9 (0)	7 (0)	2 (0.1)	0.14
Tobacco/e-cigarette smoking				
Never	13500 (65.6)	12832 (68.2)	668 (37.9)	< 0.001
Former	4567 (22.2)	3695 (19.6)	872 (49.5)	
Current	2520 (12.2)	2297 (12.2)	223 (12.6)	

SD: standard deviation; TB: tuberculosis; COPD: chronic obstructive pulmonary disease.

developed areas, such as Zona Sul), to be White, to have completed superior education or higher and to live alone; they were less likely to be currently working, to commute to work and to attend public events with 10 people or more. Comparing the two groups of participants aged 18–59 years, those with at least one comorbidity were more likely to be Pardo or Black, to have lower education (complete secondary

education or lower), to be current or former smokers, and less likely to attend public events with 10+ people.

A total of 8,780 (42.6%) participants reported ever testing for SARS-CoV-2; the proportion was lower among those aged ≥ 60 years ($p < 0.001$) (Table 3). Most of them reported a previous antibody test (75.6%) and 36.4% reported a previous PCR test. Among individuals reporting a prior SARS-CoV-2

Table 3 – Self-reported SARS-CoV-2 testing and results among study participants, Rio de Janeiro, Brazil, 2020.

	Total	18–59 Years		≥ 60 years	p-value
		No comorbidity	At least one comorbidity		
Ever tested for SARS-CoV-2 (N = 20,587)	8780 (42.6)	5073 (42.7)	3055 (44.0)	652 (37.0)	<.001
Ever performed PCR test (N = 8780)	3196 (36.4)	1861 (36.7)	1123 (36.8)	212 (32.5)	0.33
Ever performed antibody test (N = 8780)	6633 (75.6)	3848 (75.9)	2294 (75.2)	491 (75.4)	0.49
PCR result (N = 3196)					0.009
Positive	511 (16.0)	309 (16.6)	183 (16.3)	19 (9.0)	
Negative	2562 (80.2)	1484 (79.7)	887 (79.0)	191 (90.1)	
Inconclusive	47 (1.5)	25 (1.3)	21 (1.9)	1 (0.5)	
Pending result	76 (2.4)	43 (2.3)	32 (2.8)	1 (0.5)	
Antibody result (N = 6633)					0.69
Positive	664 (10.0)	381 (9.9)	240 (10.5)	43 (8.8)	
Negative	5751 (86.7)	3337 (86.7)	1978 (86.3)	436 (89)	
Inconclusive	123 (1.9)	74 (1.9)	44 (1.9)	5 (1)	
Pending result	92 (1.4)	56 (1.5)	30 (1.3)	6 (1.2)	

serology, prevalence of a positive result was 10.0%, with no difference across groups. Of those reporting prior PCR, prevalence of a positive result was 16.0%, higher among individuals aged 18–59 years compared to the those aged ≥ 60 years ($p < .009$).

Dividing the number of participants by the population estimates of each city of the Rio de Janeiro metropolitan area, residents from Rio de Janeiro city (0.26%) and Niterói (0.23%) were more frequently represented in the study sample. Within the municipality of Rio de Janeiro, residents of administrative regions with higher HDI were more frequently represented (Table 4). Respondents from higher HDI areas reported SARS-CoV-2 testing more often ($>50\%$) than those from lower HDI areas (29%) (Table 4). Self-reported overall SARS-CoV-2 prevalence (positive PCR or antibody tests) varied across regions. Among the areas with >30 volunteers, prevalence of SARS-CoV-2 by PCR was highest in areas with lower HDI.

Discussion

We have effectively reached out to a large and diverse population of individuals living in Rio de Janeiro metropolitan area through online and community recruitment strategies, indicating the willingness of the Brazilian population to register for participating in COVID 19 vaccine studies. In addition, we observed disparities in SARS-CoV-2 testing throughout different geographic regions in Rio de Janeiro metropolitan area, which may be a proxy of health disparities, including access to health services.

This study included a high proportion of white, more educated individuals from high HDI areas. However, it is worth noticing that most of the participants were unable to strictly comply to social distancing measures, therefore posing themselves at risk for SARS-CoV-2 infection. Most participants received visits at home, needed to commute to work and used public transportation (buses, trains, metro). Social distancing measures have been irregularly adopted in Brazil since the beginning of the COVID-19 epidemic.^{3,6} By opposing economy and health, the Brazilian government strategy has been to try to achieve herd immunity through contagion.²⁸ The economic crisis in the country, which had already impacted the social achievements obtained until 2013, gained a greater negative momentum in 2020 with the COVID-19 pandemic.²⁹ Our group has already described an association between socioeconomic/racial disparities and unattainability in maintaining social distancing among sexual and gender minorities.⁶ Brazilian Government's response to the pandemic has been marked by controversy about social distancing policies,³ but these policies should be considered, implemented and maintained while the population is being vaccinated.

Our findings indicate that older individuals (aged ≥ 60 years) were less exposed to COVID-19 in terms of social distancing and performed less SARS-CoV-2 tests than individuals aged 18–59 years. Among the later group, participants with comorbidities were less exposed to COVID-19 and more frequently tested for SARS-CoV-2. This is in accordance with a previous study conducted in Rio Grande Sul, Brazil,³⁰ and is relevant, as older individuals and persons with comorbidities are more likely to develop severe COVID-19 and death.³¹

Conversely, results from a recent study showed that Brazilian individuals who are male, older, with lower education levels, and living in less developed regions had lower knowledge about COVID-19,³² which may expose them to increased risk of infection.

Less than half (42.6%) of participants reported ever testing for SARS-CoV-2. SARS-CoV-2 prevalence according to self-reported PCR and antibody positive test results were 16.0% and 10.0%, respectively. These results indicate that individuals somehow affected by COVID-19 pandemic may be more interested in COVID-19 vaccines, as SARS-CoV-2 testing is more frequently performed by individuals who are symptomatic or have more severe illnesses.³³ SARS-CoV-2 antibody prevalence in Rio de Janeiro city was estimated at 2.4% [95% confidence interval – 95% CI: 0.7–5.6] during May 14–21, 2020 and 7.5% [95% CI: 4.5–11.7] during June 4–7, 2020.³⁴ Increase in antibody prevalence was expected as our study was conducted some months after the last assessment (from September to December 2020). Nevertheless, our results should be analyzed with caution as we collected self-reported data and diagnostic methods and results may be overestimated.

This study observed disparities in SARS-CoV-2 testing across different regions of Rio de Janeiro metropolitan area. Overall, more individuals from higher HDI areas reported ever being tested for SARS-CoV-2, indicating disparities in access to health care, as tests are not widely available at the Brazilian Public Health System (SUS) and focus on symptomatic individuals. Individuals living in higher HDI areas may have access to tests through private laboratories and clinics (less prone to shortage in testing). Despite universal susceptibility to COVID-19, populations with increased social vulnerability in Brazil are more and unequally affected by COVID-19,³⁵ indicating that policies to prioritize low income individuals need to be considered and urgently implemented.

Important considerations about the online registry should be noted. Fiocruz website has shown to be highly effective for recruitment, indicating population trust in scientific research conducted by this centenary institution and its importance in providing scientific information. Our study population mostly comprised individuals aged 18–59 years, despite our efforts to increase recruitment of older individuals through social media advertisements. Community activities were important to increase recruitment of individuals from lower HDI areas, although the majority of individuals recruited were from higher HDI areas. Even though internet access has increased in Brazil among all social strata,³⁶ structural social factors such as access to education may have prevented individuals from lower HDI areas to access and complete the questionnaire.

Although this study was performed in a single geographic location and included only individuals willing to participate in clinical trials for COVID-19 vaccines, some characteristics of our sample were similar to the general population, suggesting that our results may have external validity. Prevalence of comorbidities such as hypertension (13.3%), diabetes (5.3%), cardiovascular diseases (2.6%) and cancer (1.0%) were similar to those estimated in the most recent national survey (data collected in November 2020)³⁷ and the prevalence of tobacco smoking was similar to national data from 2019 (12.6%).³⁸ Moreover, we present data from a large sample of over 20,000

Table 4 – Self-reported SARS-CoV-2 testing and results among study participants according to municipality and Rio de Janeiro administrative area. Rio de Janeiro, Brazil, 2020.

	Pop.	% Pop.	Ever tested			PCR			Antibody		
			N	Yes	%	N	+	%	N	+	%
Rio de Janeiro city	6,747,815	0.26	17380	7444	42.8	2817	430	15.3	5686	549	9.7
HDI > 0.9											
Lagoa	163,139	0.35	570	338	59.3	176	15	8.5	261	15	5.7
Barra da Tijuca	394,037	0.28	1085	595	54.8	280	34	12.1	439	31	7.1
Copacabana	161,201	0.49	790	408	51.6	161	16	9.9	323	21	6.5
Botafogo	240,344	0.68	1628	798	49.0	322	21	6.5	641	36	5.6
Tijuca	182,413	0.64	1171	537	45.9	194	24	12.4	432	38	8.8
Vila Isabel	191,740	0.49	934	414	44.3	151	24	15.9	319	33	10.3
HDI = 0.850-0.899											
Centro	42,621	0.77	328	144	43.9	44	7	15.9	114	13	11.4
Santa Teresa	40,765	0.34	138	59	42.8	21	5	23.8	41	2	4.9
Méier	397,263	0.36	1418	582	41.0	204	41	20.1	442	49	11.1
Irajá	202,941	0.32	649	229	35.3	84	22	26.2	169	23	13.6
Ilha do Governador	213,388	0.35	744	259	34.8	117	17	14.5	196	20	10.2
HDI = 0.800-0.849											
Ramos	155,222	0.35	538	230	42.8	78	15	19.2	170	27	15.9
Jacarepaguá	648,484	0.25	1622	683	42.1	265	52	19.6	514	51	9.9
Inhaúma	137,086	0.27	370	152	41.1	50	9	18.0	109	17	15.6
Rio Comprido	82,892	0.25	211	84	39.8	28	7	25.0	63	5	7.9
Madureira	370,355	0.17	648	228	35.2	75	17	22.7	172	22	12.8
Realengo	245,851	0.12	303	105	34.7	40	3	7.5	74	6	8.1
São Critóvão	95,199	0.23	216	74	34.3	33	9	27.3	49	12	24.5
Penha	187,575	0.19	359	120	33.4	39	8	20.5	96	8	8.3
Vigário Geral	136,805	0.15	211	60	28.4	15	4	26.7	49	6	12.2
Anchieta	161,052	0.13	208	55	26.4	20	3	15.0	39	8	20.5
Ilha de Paqueta	3317	0.36	12	1	8.3	1	0	0.0	0	0	0.0
HDI = 0.750-799											
Cidade de Deus	35,409	0.03	12	6	50.0	1	0	0.0	6	1	16.7
Campo Grande	584,628	0.13	771	256	33.2	88	21	23.9	189	21	11.1
Bangu	433,586	0.08	352	114	32.4	38	12	31.6	92	17	18.5
Zona Portuária	55,070	0.12	64	20	31.2	8	0	0.0	13	1	7.7
Pavuna	217,470	0.08	175	47	26.9	19	5	26.3	28	9	32.1
HDI < 0.750											
Complexo do Alemão	72,177	0.01	6	5	83.3	1	0	0.0	5	0	0.0
Maré	141,535	0.06	79	32	40.5	14	4	28.6	22	6	27.3
Jacarezinho	38,856	0.05	18	6	33.3	1	0	0.0	4	0	0.0
Santa Cruz	410,726	0.05	218	64	29.4	18	4	22.2	48	8	16.7
Guaratiba	139,262	0.07	103	30	29.1	12	2	16.7	18	3	16.7
Rocinha	78,951	0.04	30	8	26.7	2	0	0.0	5	1	20.0
Metropolitan cities											
HDI = 0.800-0.849											
Niterói	515,317	0.23	1206	593	49.2	199	26	13.1	465	31	6.7
HDI = 0.750-0.799											
Rio Bonito	60,573	0.05	32	14	43.8	4	0	0.0	11	0	0.0
Maricá	164,504	0.08	138	56	40.6	28	3	10.7	43	4	9.3
Nilópolis	162,693	0.10	168	65	38.7	20	5	25.0	46	7	15.2
Cachoeiras de Macacu	59,303	0.04	22	8	36.4	2	0	0.0	8	1	12.5
Petrópolis	306,678	0.05	144	78	54.2	29	6	20.7	64	6	9.4
HDI = 0.700-0.749											
São Gonçalo	1,091,737	0.06	618	236	38.2	54	10	18.5	179	20	11.2
Mesquita	176,569	0.07	122	55	45.1	18	7	38.9	34	8	23.5
Paracambi	52,683	0.03	16	6	37.5	1	0	0.0	4	0	0.0
São João de Meriti	472,906	0.07	314	129	41.1	38	10	26.3	89	15	16.9
Itaguaí	134,819	0.04	57	17	29.8	3	1	33.3	11	1	9.1
Nova Iguaçu	823,302	0.06	493	220	44.6	64	16	25.0	161	20	12.4
Seropédica	83,092	0.11	95	26	27.4	5	0	0.0	14	1	7.1
Duque de Caxias	924,624	0.07	659	301	45.7	85	19	22.4	202	24	11.9
Magé	246,433	0.04	104	25	24.0	6	1	16.7	16	2	12.5
Itaboraí	242,543	0.04	94	30	31.9	6	1	16.7	23	2	8.7
Guapimirim	61,388	0.05	29	16	55.2	6	1	16.7	10	2	20.0
HDI < 0.700											
Belford Roxo	513,118	0.04	193	108	56.0	18	3	16.7	79	8	10.1
Queimados	151,335	0.05	71	26	36.6	7	1	14.3	21	2	9.5
Japeri	105,548	0.02	21	7	33.3	3	0	0.0	6	0	0.0
Tanguá	34,610	0.03	10	3	30.0	0	0	0.0	2	0	0.0
Total	13,131,590	0.16	20587	8780	42.6	3196	511	16.0	6633	664	10.0

¹Estimated population of each city and region in 2020.

²%pop: percentage of individuals who tested for SARS-CoV-2 considering the estimated population of each city and region in 2020. HDI: Human Development Index.

participants living in a large, diverse metropolitan area in Brazil, which indicates socioeconomic disparities in access to health care.

This study has limitations. First, web-based studies are not probabilistic sampling strategies, precluding the generalization of the findings to all individuals living in Rio de Janeiro metropolitan area or other metropolitan areas or regions in Brazil. Moreover, our findings are based on those who have access to mobile phones and internet connection. Nevertheless, recent data show that 79% of Brazilians have access to internet connection³⁶ and 85% have mobile phones.³⁹ All collected data were self-reported by participants and may be subject to bias, although individuals tend to be more honest through web-based surveys, reducing social desirability bias.⁴⁰ Information on social distancing and mobility may be subject to recall bias or social expectation bias, so these results may be overestimated.

In conclusion, interest to participate in COVID-19 vaccines clinical trials in Rio de Janeiro metropolitan area, Brazil was very high. A large number of individuals with diverse sociodemographic, economic and clinical backgrounds were successfully registered for potential participation in clinical trials through an online platform. Our procedures to register participants interested on clinical trials for COVID-19 vaccines could be useful to other research centers in Brazil or in other countries to recruit individuals to clinical trials in general, not only to COVID-19 vaccine trials.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

TST acknowledges funding from the National Council of Technological (CNPq, #28/2018). BG acknowledges funding from the National Council of Technological and Scientific Development and the Research Funding Agency of the State of Rio de Janeiro (Programa Cientista do Nosso Estado; Edital N.º 03/2018).

REFERENCES

- World Health Organization (WHO). WHO Director-General's Opening Remarks at the Media Briefing on COVID-19 - 11. World Health Organization (WHO); 2020. March Available at: <https://www.who.int/dg/speeches/detail/> [accessed May 18, 2021].
- Brasil, Governo Federal. Paineis Coronavírus. Brasil, Governo Federal Available at: <https://covid.saude.gov.br> [accessed August 11, 2021].
- The Lancet COVID-19 in Brazil: "so what?". The Lancet. 2020;395(10235):1461. [https://doi.org/10.1016/S0140-6736\(20\)31095-3](https://doi.org/10.1016/S0140-6736(20)31095-3).
- Folha de S. Paulo. Isolamento Atinge Níveis Mais Baixos Desde Início da Pandemia, Aponta Datafolha. Folha de S. Paulo; 2021. Available at <https://www1.folha.uol.com.br/equilibrioesaude/2021/05/isolamento-atinge-niveis-mais-baixos-desde-inicio-da-pandemia-aponta-datafolha.shtml> [accessed May 18, 2021].
- Dall'Alba R, Rocha CF, de Pinho Silveira R, da Silva Costa Dresch L, Vieira LA, Germanò MA. COVID-19 in Brazil: far beyond biopolitics. The Lancet. 2021;397(10274):579–80. [https://doi.org/10.1016/S0140-6736\(21\)00202-6](https://doi.org/10.1016/S0140-6736(21)00202-6).
- Torres TS, Hoagland B, Bezerra DRB, et al. Impact of COVID-19 pandemic on sexual minority populations in Brazil: an analysis of social/racial disparities in maintaining social distancing and a description of sexual behavior. AIDS Behav. 2021;25(1):73–84. <https://doi.org/10.1007/s10461-020-02984-1>.
- United Nations Economic Commission for Latin America and the Caribbean Social Panorama of Latin America 2019. Available at: <https://www.cepal.org/en/publications/44989-social-panorama-latin-america-2019> [accessed June 4, 2020].
- Worldometer. World population Available at: <https://www.worldometers.info/world-population/population-by-country> [accessed May 18, 2021].
- Bilal U, Alazraqui M, Caiaffa WT, et al. Inequalities in life expectancy in six large Latin American cities from the SALURBAL study: an ecological analysis. Lancet Planet Health. 2019;3(12):e503–10. [https://doi.org/10.1016/S2542-5196\(19\)30235-9](https://doi.org/10.1016/S2542-5196(19)30235-9).
- Baqui P, Bica I, Marra V, Ercole A, van der Schaar M. Ethnic and regional variations in hospital mortality from COVID-19 in Brazil: a cross-sectional observational study. Lancet Glob Health. 2020;8(8):e1018–26. [https://doi.org/10.1016/S2214-109X\(20\)30285-0](https://doi.org/10.1016/S2214-109X(20)30285-0).
- World Health Organization (WHO). Framework for Decision-Making: Implementation of Mass Vaccination Campaigns in the Context of COVID-19. World Health Organization (WHO); 2021. Available at https://apps.who.int/iris/bitstream/handle/10665/332159/WHO-2019-nCoV-Framework_Mass_Vaccination-2020.1-eng.pdf [accessed May 18, 2021].
- Voysey M, Clemens SAC, Madhi SA, et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. The Lancet. 2021;397(10269):99–111. [https://doi.org/10.1016/S0140-6736\(20\)32661-1](https://doi.org/10.1016/S0140-6736(20)32661-1).
- Bio-Manguinhos. Saiba mais sobre a vacina COVID-19. Available at: <https://www.bio.fiocruz.br/index.php/br/perguntas-e-respostas-a-vacina> [accessed April 13, 2021].
- Polack FP, Thomas SJ, Kitchin N, et al. Safety and efficacy of the BNT162b2 mRNA Covid-19 Vaccine. N Engl J Med. 2020;383(27):2603–15. <https://doi.org/10.1056/NEJMoa2034577>.
- Brasil, Ministério da Saúde. O Que Muda Com o Registro de Uma Vacina? Brasil, Ministério da Saúde; 2021. Available at <https://www.gov.br/anvisa/pt-br/assuntos/noticias-anvisa/2021/o-que-muda-com-o-registro-de-uma-vacina> [accessed May 18, 2021].
- Zhang Y, Zeng G, Pan H, et al. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. Lancet Infect Dis. 2021;21(2):181–92. [https://doi.org/10.1016/S1473-3099\(20\)30843-4](https://doi.org/10.1016/S1473-3099(20)30843-4).
- Instituto Butantan. Bula CoronaVac. Available at: <https://vacinacovid.butantan.gov.br/bulas> [accessed May 18, 2021].
- Sadoff J, Le Gars M, Shukarev G, et al. Interim results of a Phase 1–2a Trial of Ad26.COV2.S Covid-19 Vaccine. N Engl J Med. 2021. <https://doi.org/10.1056/NEJMoa2034201>.
- Globo G1. Mapa da vacinação contra Covid-19 no Brasil. Available at: <https://especiais.g1.globo.com/bemestar/vacina/2021/mapa-brasil-vacina-covid/> [accessed August 11, 2021].
- Shahcheraghi SH, Ayatollahi J, Aljabali AA, et al. An overview of vaccine development for COVID-19. Ther Deliv. 2021;12(3):235–44. <https://doi.org/10.4155/tde-2020-0129>.

21. Instituto Nacional de Infectologia Evandro Chagas INI-Fiocruz. Pesquisa no INI-Fiocruz. Instituto Nacional de Infectologia Evandro Chagas INI-Fiocruz; 2021. Available at <https://www.ini.fiocruz.br/pesquisa> [accessed April 14, 2021].
22. Eysenbach G. Improving the quality of web surveys: the checklist for reporting results of internet E-surveys (CHERRIES). *J Med Internet Res.* 2004;6(3):e34. <https://doi.org/10.2196/jmir.6.3.e34>.
23. IBGE. Cidades e Estados. Available at: <https://www.ibge.gov.br/cidades-e-estados/rj/rio-de-janeiro.html> [accessed April 7, 2021].
24. Decreto n.o 3158. Available at: <http://www2.rio.rj.gov.br/smu/buscafacil/Arquivos/PDF/D3158M.PDF> [accessed April 7, 2021].
25. Instituto Pereira Passos. População Residente e Estimada - Brasil, Estado do Rio de Janeiro e Município do Rio de Janeiro e Regiões Administrativas (RA) - 2000/2010/2013-2016/2020. Available at: <https://www.data.rio/datasets/ba877d53302346eca990a47c99e15f74> [accessed April 7, 2021].
26. Instituto Pereira Passos. Índice de Desenvolvimento Humano (IDH) Geral por Regiões Administrativas (RA) do Município do Rio de Janeiro. Instituto Pereira Passos. 2018. Available at: <https://www.data.rio/datasets/6d43a854cae84bc692074d0af4e922d9>. [accessed August 10, 2021].
27. The R project for statistical computing. Available at: <https://www.r-project.org/> [accessed May 26, 2020].
28. Milani CRS, Nery T. Brazil's Catastrophic Response to Covid-19: The International–Domestic Nexus. *Berkeley Review of Latin American Studies*; 2021 [In Press].
29. Rede PENSSAN. VIGISAN Inquérito Nacional sobre Insegurança Alimentar no Contexto da Pandemia da Covid-19 no Brasil. Available at: <http://www.olheparaafome.com.br/> [accessed May 17, 2021].
30. Barros AJD, Victora CG, Menezes AMB, et al. Social distancing patterns in nine municipalities of Rio Grande do Sul, Brazil: the Epicovid19/RS study. *Rev Saúde Pública.* 2020;54:75. <https://doi.org/10.11606/s1518-8787.2020054002810>.
31. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020;584(7821):430–6. <https://doi.org/10.1038/s41586-020-2521-4>.
32. Guimarães VHA, de Oliveira-Leandro M, Cassiano C, et al. Knowledge about COVID-19 in Brazil: cross-sectional web-based study. *JMIR Public Health Surveill.* 2021;7(1):e24756. <https://doi.org/10.2196/24756>.
33. França EB, Ishitani LH, Teixeira RA, et al. Óbitos por COVID-19 no Brasil: quantos e quais estamos identificando? *Rev Bras Epidemiol.* 2020;23:e200053. <https://doi.org/10.1590/1980-549720200053>.
34. Hallal PC, Hartwig FP, Horta BL, et al. SARS-CoV-2 antibody prevalence in Brazil: results from two successive nationwide serological household surveys. *Lancet Glob Health.* 2020;8(11):e1390–8. [https://doi.org/10.1016/S2214-109X\(20\)30387-9](https://doi.org/10.1016/S2214-109X(20)30387-9).
35. Raymundo CE, Oliveira MC, Eleuterio T de A, et al. Spatial analysis of COVID-19 incidence and the sociodemographic context in Brazil. *PLoS ONE.* 2021;16(3):e0247794. <https://doi.org/10.1371/journal.pone.0247794>.
36. Comitê Gestor da Internet no Brasil (CGIBR). C2A - USUÁRIOS DE INTERNET - INDICADOR AMPLIADO 2019. Available at: <https://cetic.br/pt/tics/domicilios/2019/individuos/C2A/> [accessed May 17, 2021].
37. Instituto Brasileiro de Geografia e Estatística (IBGE). PNAD COVID-19 mensal - Novembro 2020. Available at: <https://cidades.ibge.gov.br/brasil/pesquisa/10095/0> [accessed July 5, 2021].
38. Instituto Nacional do Câncer (INCA). Dados e números da prevalência do tabagismo. Available at: <https://www.inca.gov.br/observatorio-da-politica-nacional-de-controle-do-tabaco/dados-e-numeros-prevalencia-tabagismo> [accessed July 5, 2021].
39. Comitê Gestor da Internet no Brasil (CGIBR). J2 - Indivíduos que possuem telefone celular 2019. Available at: <https://cetic.br/pt/tics/domicilios/2019/individuos/J2/> [accessed May 17, 2021].
40. Heerwegh D. Mode differences between face-to-face and web surveys: an experimental investigation of data quality and social desirability effects. *Int J Public Opin Res.* 2009;21(1):111–21. <https://doi.org/10.1093/ijpor/edn054>.